

To: _____

From: _____

The People's Notice to Federal Congress

I, _____, one of the People of the 50 republican states, do provide you with due notice of danger to the People, that you may provide due care;

I, one of the People, wish to give you this notice with attached evidence of findings that may bring harm to the People of America who may take the Covid 19 Shot;

As one of the People, I wish to declare that many of the People have come together to consult for the common good and found that there are Title 42 Cooperative programs where money is being made between the Federal, State, and Political Subdivision workers and other private actors which may present a conflict of interest where unlawful jurisdiction has been extended over the free People of the 50 Republican States;

Please take notice that as one of the People, I wish for you to show the People all Covid related funds that are being paid out in any partnership, with a full accounting of where all funds were distributed;

Please take further notice that if there are any private interests that you have in any of these programs, that you are acting with unclean hands and giving influence to the State and Political Subdivisions in actions that are nefarious and harmful to the People;

As one of the People, I wish for you to show where in Article 1 Section 8 you were granted authority, by the People, to make rules for the People, and if you believe that use of some commerce clause trick allows for you to gain jurisdiction over the free people, please show constitutional authority and where you gave full disclosure to the People;

Please take further notice that I wish for you to show constitutional authority you have been granted authority to limit the liberty of the People in a republic (See Constitutional Evidence below):

Kentucky Constitution Bill of Rights Text of Section 2:

Absolute and arbitrary power over the lives, liberty and property of freemen exists nowhere in a republic, not even in the largest majority.

Please take notice that all 50 states are guaranteed a republican form of government;

United States Constitution Article 4 Section 4

Article IV

Section 4

The United States shall guarantee to every State in this Union a Republican Form of Government, and shall protect each of them against Invasion; and on Application of the Legislature, or of the Executive (when the Legislature cannot be convened) against domestic Violence.

Please take notice that as one of the People, I realize you are not allowed to remove my right to life, liberty and property, no matter if all Legislative Officials voted, nor can the President, make a rule that would infringe on the People's freedom (Please see evidence below):

Miranda v. Arizona, 384 U.S. 436 (1966)

Where rights secured by the Constitution are involved, there can be no rulemaking or legislation which would abrogate them.

Please take notice that the roles have somehow been turned about and where you may believe you lead us, in a republic you are simply a servant to act as a trustee. Therefore, all the conflicts of interest and failures to follow the limitations outlined in the 51 constitutions are now causing distress and harm to the People. Therefore, we have returned to our rightful understanding and are ready to instruct our servants as seen below in our constitutions:

Michigan Constitution Declaration of Rights:

Text of Section 3:

Assembly, Consultation, Instruction, Petition

The people have the right peaceably to assemble, to consult for the common good, to **instruct their representatives** and to petition the government for redress of grievances.[Emphasis by Highlight Added]

Idaho Constitution Text of Section 10:

Right of Assembly

The people shall have the right to assemble in a peaceable manner, to consult for their common good; to **instruct their representatives**, and to petition the legislature for the redress of grievances.

Texas Constitution Bill of Rights:

Text of Section 2:

Inherent Political Power; Republican Form of Government

All political power is inherent in the people, and all free governments are founded on their authority, and instituted for their benefit. The faith of the people of Texas stands pledged to the preservation of a **republican form of government**, and, **subject to this limitation only, they have at all times the inalienable right to alter, reform or abolish their government in such manner as they may think expedient.** [Emphasis by Highlight Added]

Please take further notice that our forefathers show us that the People have the power to reform, alter, or abolish government WHENEVER THEY MAY THINK IT EXPEDIENT(not waiting for the next general election);

Please take further notice that, if we have a right to reform, alter or abolish government or constitutions for our safety, it would be against your oath and office to deny, infringe, or interfere with this right. Therefore, I wish for you to immediately remove all mandates and educate every one of your agents that they do not have any authority over the free people and that slavery was abolished long ago! If you believe that you have been given constitutional authority to deny the private rights of the People, please respond within 10 days with those provisions, rebutting this notice, point by point in the form of a sworn affidavit. If you fail to respond or take decisive action to relieve the People, based on the instructions above, you agree that you are knowingly, maliciously, and with full intent, denying the rights of the People and ignoring the oath you made as a purposeful trespass. The People also wish to have direct access to thier federal grand juries to correct any government error and not have any Bar Association member controlling the People's best interest and control. This notice is sent to you in peace and the love of Christ, that you may provide due care to those who have all political power, the People.

Autograph:

Date:

To:

From: Dr. Jane Ruby

Affidavit of Findings

Affiant, Jane Ruby, PhD, EdD, MS, one of the People, Sui Juris, in this Court of Record am presenting the following claims:


1. Affiant has found the following evidence of harmful items to the People of America while engaging in research: Polyethylene glycol in the Pfizer-BioNTech Covid 19 Vaccine is toxic to human beings and known to cause anaphylaxis and cardiac arrest.
2. Presence in Pfizer BioNTech Covid 19 Vaccine: See Exhibit 1: (Pfizer Prescriber Information)
3. Effects on Humans: See Exhibit 2: (Sellaturay P, Nasser S, Ewan P. Polyethylene Glycol-Induced Systemic Allergic Reactions (Anaphylaxis). J Allergy Clin Immunol Pract. 2021 Feb;9(2):670-675.)

Affiant declares that the paramount duty of government is the safety and protection of the People and their property, therefore Affiant demands immediate investigation into these matters immediately and any potential funds made from any federal program dealing with Covid to be presented to Affiant as it may be a potential conflict of interest.

Verification

I hereby declare, certify, and state, pursuant to the penalties of perjury under the laws of the United States of America, and by the provisions of 28 USC § 1746 that all of the above and foregoing representations are true and correct to the best of my knowledge, information and belief.

Executed in Palm Beach, Florida) on this 13th day of December in the Year of Our Lord Two Thousand and Twenty-One.

 Autograph of Affiant:

Notary as JURAT CERTIFICATE


Florida State} Palm Beach County}

On this 13 day of December, 2021 (date) before me, _____, a Notary Public, personally appeared Dr. Jane Ruby, Name of Affiant, who proved to me on the basis of satisfactory evidence to be the woman whose name is subscribed to the within instrument and acknowledged to me that she executed the same in her authorized capacity, and that by her autograph(s) on the instrument the woman executed, the instrument.

I certify under PENALTY OF PERJURY under the lawful laws of the State of Florida and that the foregoing paragraph is true and correct.

WITNESS my hand and official seal.

Signature of Notary / Jurat _____



**FACT SHEET FOR HEALTHCARE PROVIDERS ADMINISTERING VACCINE
(VACCINATION PROVIDERS)**

**EMERGENCY USE AUTHORIZATION (EUA) OF
THE PFIZER-BIONTECH COVID-19 VACCINE TO PREVENT CORONAVIRUS
DISEASE 2019 (COVID-19)**

**FOR 12 YEARS OF AGE AND OLDER
DILUTE BEFORE USE**

The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) to permit the emergency use of the unapproved product, Pfizer-BioNTech COVID-19 Vaccine, for active immunization to prevent COVID-19 in individuals 5 years of age and older.

There are 2 formulations of Pfizer-BioNTech COVID-19 Vaccine authorized for use in individuals 12 years of age and older:

**The formulation supplied in a multiple dose vial with a purple cap
MUST BE DILUTED PRIOR TO USE.**

**The formulation supplied in a multiple dose vial with a gray cap and
label with a gray border IS NOT DILUTED PRIOR TO USE.**

**This Fact Sheet pertains only to Pfizer-BioNTech COVID-19 Vaccine supplied
in a multiple dose vial with a purple cap, which is authorized for use in
individuals 12 years of age and older and MUST BE DILUTED PRIOR TO USE.**

**Pfizer-BioNTech COVID-19 Vaccine supplied in a multiple dose vial with a
purple cap is authorized for use to provide:**

- **a 2-dose primary series to individuals 12 years of age and older;**
- **a third primary series dose to individuals 12 years of age and older
who have been determined to have certain kinds of
immunocompromise;**
- **a single booster dose to individuals 16 years of age and older who
have completed a primary series with Pfizer-BioNTech COVID-19
Vaccine or COMIRNATY; and**
- **a single booster dose to individuals 18 years of age and older who
have completed primary vaccination with a different authorized
COVID-19 vaccine. The dosing interval for the heterologous booster
dose is the same as that authorized for a booster dose of the vaccine
used for primary vaccination.**

**COMIRNATY (COVID-19 Vaccine, mRNA) is an FDA-approved COVID-19
vaccine made by Pfizer for BioNTech that is indicated for active immunization
to prevent COVID-19 in individuals 16 years of age and older. It is approved
for use as a 2-dose primary series for the prevention of COVID-19 in**

individuals 16 years of age and older. It is also authorized for emergency use to provide:

- a 2-dose primary series to individuals 12 through 15 years of age;
- a third primary series dose to individuals 12 years of age and older who have been determined to have certain kinds of immunocompromise;
- a single booster dose to individuals 16 years of age and older who have completed a primary series with Pfizer-BioNTech COVID-19 Vaccine or COMIRNATY; and
- a single booster dose to individuals 18 years of age and older who have completed primary vaccination with a different authorized COVID-19 vaccine. The dosing interval for the heterologous booster dose is the same as that authorized for a booster dose of the vaccine used for primary vaccination.

The FDA-approved COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine for individuals 12 years of age and older when prepared according to their respective instructions for use can be used interchangeably.¹

COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine intended for individuals 12 years of age and older should not be used for individuals 5 through 11 years of age because of the potential for vaccine administration errors, including dosing errors.²

SUMMARY OF INSTRUCTIONS FOR COVID-19 VACCINATION PROVIDERS

Vaccination providers enrolled in the federal COVID-19 Vaccination Program must report all vaccine administration errors, all serious adverse events, cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and cases of COVID-19 that result in hospitalization or death following administration of Pfizer-BioNTech COVID-19 Vaccine. See “MANDATORY REQUIREMENTS FOR PFIZER-BIONTECH COVID-19 VACCINE ADMINISTRATION UNDER EMERGENCY USE AUTHORIZATION” for reporting requirements.

¹ When prepared according to their respective instructions for use, the FDA-approved COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine for individuals 12 years of age and older can be used interchangeably without presenting any safety or effectiveness concerns.

² Notwithstanding the age limitations for use of the different formulations and presentations described above, individuals who will turn from 11 years to 12 years of age between their first and second dose in the primary regimen may receive, for either dose, either: (1) the Pfizer-BioNTech COVID-19 Vaccine formulation authorized for use in individuals 5 through 11 years of age (each 0.2 mL dose containing 10 mcg modRNA) (orange cap); or (2) COMIRNATY or one of the Pfizer-BioNTech COVID-19 Vaccine formulations authorized for use in individuals 12 years of age and older (each 0.3 mL dose containing 30 mcg modRNA) (gray and purple cap).

The Pfizer-BioNTech COVID-19 Vaccine is a suspension for intramuscular injection.

Primary Series

The Pfizer-BioNTech COVID-19 Vaccine is administered as a primary series of 2 doses (0.3 mL each) 3 weeks apart in individuals 12 years of age or older.

A third primary series dose of the Pfizer-BioNTech COVID-19 Vaccine (0.3 mL) at least 28 days following the second dose is authorized for administration to individuals at least 12 years of age who have undergone solid organ transplantation, or who are diagnosed with conditions that are considered to have an equivalent level of immunocompromise.

Booster Dose

A single Pfizer-BioNTech COVID-19 Vaccine booster dose (0.3 mL) may be administered at least 6 months after completing a primary series of the Pfizer-BioNTech COVID-19 Vaccine or COMIRNATY to individuals 16 years of age and older.

A single booster dose of the Pfizer-BioNTech COVID-19 Vaccine may be administered to individuals 18 years of age and older as a heterologous booster dose following completion of primary vaccination with another authorized COVID-19 vaccine. The dosing interval for the heterologous booster dose is the same as that authorized for a booster dose of the vaccine used for primary vaccination.

See this Fact Sheet for instructions for preparation and administration. This Fact Sheet may have been updated. For the most recent Fact Sheet, please see www.cvdvaccine.com.

For information on clinical trials that are testing the use of the Pfizer-BioNTech COVID-19 Vaccine for active immunization against COVID-19, please see www.clinicaltrials.gov.

DESCRIPTION OF COVID-19

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by the novel coronavirus, SARS-CoV-2, that appeared in late 2019. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have reported a wide range of symptoms, ranging from mild symptoms to severe illness. Symptoms may appear 2 to 14 days after exposure to the virus. Symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

DOSAGE AND ADMINISTRATION

The storage, preparation, and administration information in this Fact Sheet apply to the Pfizer-BioNTech COVID-19 Vaccine for individuals 12 years of age and older, which is supplied in a multiple dose vial with a purple cap and **MUST BE DILUTED** before use.

Pfizer-BioNTech COVID-19 Vaccine, Multiple Dose Vial with Purple Cap

Age Range	Dilution Information	Doses Per Vial After Dilution	Dose Volume
12 years and older	Dilute with 1.8 mL sterile 0.9% Sodium Chloride Injection, USP prior to use	6	0.3 mL

Storage and Handling

During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.

Do not refreeze thawed vials.

Frozen Vials Prior to Use

Cartons of Pfizer-BioNTech COVID-19 Vaccine multiple dose vials with purple caps arrive in thermal containers with dry ice. Once received, remove the vial cartons immediately from the thermal container and preferably store in an ultra-low temperature freezer between -90°C to -60°C (-130°F to -76°F) until the expiry date printed on the label. This information in the package insert supersedes the storage conditions printed on the vial cartons.

Cartons and vials of Pfizer-BioNTech COVID-19 Vaccine supplied in multiple dose vials with purple caps with an expiry date of August 2021 through February 2022 printed on the label may remain in use for 3 months beyond the printed date as long as approved storage conditions between -90°C to -60°C (-130°F to -76°F) have been maintained. Updated expiry dates are shown below.

<u>Printed Expiry Date</u>		<u>Updated Expiry Date</u>
August 2021	→	November 2021
September 2021	→	December 2021
October 2021	→	January 2022
November 2021	→	February 2022
December 2021	→	March 2022
January 2022	→	April 2022
February 2022	→	May 2022

If not stored between -90°C to -60°C (-130°F to -76°F), vials may be stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks. Vials must be kept frozen and protected

from light until ready to use. Vials stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks may be returned one time to the recommended storage condition of -90°C to -60°C (-130°F to -76°F). Total cumulative time the vials are stored at -25°C to -15°C (-13°F to 5°F) should be tracked and should not exceed 2 weeks.

If an ultra-low temperature freezer is not available, the thermal container in which the Pfizer-BioNTech COVID-19 Vaccine arrives may be used as temporary storage when consistently re-filled to the top of the container with dry ice. Refer to the re-icing guidelines packed in the original thermal container for instructions regarding the use of the thermal container for temporary storage. The thermal container maintains a temperature range of -90°C to -60°C (-130°F to -76°F). Storage of the vials between -96°C to -60°C (-141°F to -76°F) is not considered an excursion from the recommended storage condition.

Transportation of Frozen Vials

If local redistribution is needed and full cartons containing vials cannot be transported at -90°C to -60°C (-130°F to -76°F), vials may be transported at -25°C to -15°C (-13°F to 5°F). Any hours used for transport at -25°C to -15°C (-13°F to 5°F) count against the 2-week limit for storage at -25°C to -15°C (-13°F to 5°F). Frozen vials transported at -25°C to -15°C (-13°F to 5°F) may be returned one time to the recommended storage condition of -90°C to -60°C (-130°F to -76°F).

Thawed Vials Before Dilution

Thawed Under Refrigeration

Thaw and then store undiluted vials in the refrigerator [2°C to 8°C (35°F to 46°F)] for up to 1 month. A carton of 25 vials or 195 vials may take up to 2 or 3 hours, respectively, to thaw in the refrigerator, whereas a fewer number of vials will thaw in less time.

Thawed at Room Temperature

For immediate use, thaw undiluted vials at room temperature [up to 25°C (77°F)] for 30 minutes. Thawed vials can be handled in room light conditions. Vials must reach room temperature before dilution.

Undiluted vials may be stored at room temperature for no more than 2 hours.

Transportation of Thawed Vials

Available data support transportation of one or more thawed vials at 2°C to 8°C (35°F to 46°F) for up to 12 hours.

Vials After Dilution

- After dilution, store vials between 2°C to 25°C (35°F to 77°F) and use within 6 hours from the time of dilution.
- During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.
- Any vaccine remaining in vials must be discarded after 6 hours.
- Do not refreeze.

Dosing and Schedule

Primary Series

The Pfizer-BioNTech COVID-19 Vaccine is administered intramuscularly as a primary series of 2 doses (0.3 mL each) 3 weeks apart to individuals 12 years of age and older.

A third primary series dose of the Pfizer-BioNTech COVID-19 Vaccine (0.3 mL) at least 28 days following the second dose is authorized for administration to individuals at least 12 years of age who have undergone solid organ transplantation, or who are diagnosed with conditions that are considered to have an equivalent level of immunocompromise.

Booster Dose

A single Pfizer-BioNTech COVID-19 Vaccine booster dose (0.3 mL) may be administered at least 6 months after completing a primary series of the Pfizer-BioNTech COVID-19 Vaccine or COMIRNATY to individuals 16 years of age and older.

A single booster dose of the Pfizer-BioNTech COVID-19 Vaccine may be administered to individuals 18 years of age and older as a heterologous booster dose following completion of primary vaccination with another authorized COVID-19 vaccine. The dosing interval for the heterologous booster dose is the same as that authorized for a booster dose of the vaccine used for primary vaccination.

The FDA-approved COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine for individuals 12 years of age and older when prepared according to their respective instructions for use, can be used interchangeably.

COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine intended for individuals 12 years of age and older should not be used for individuals 5 through 11 years of age because of the potential for vaccine administration errors, including dosing errors.

Dose Preparation

Each vial **MUST BE DILUTED** before administering the vaccine.


Prior to Dilution



- The Pfizer-BioNTech COVID-19 Vaccine multiple dose vial with a purple cap contains a volume of 0.45 mL and is supplied as a frozen suspension that does not contain preservative.
- Each vial must be thawed before dilution.
 - Vials may be thawed in the refrigerator [2°C to 8°C (35°F to 46°F)] or at room temperature [up to 25°C (77°F)] (*see Storage and Handling*).
 - Refer to thawing instructions in the panels below.

Dilution

Dilute the vial contents using 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP (not provided) to form the Pfizer-BioNTech COVID-19 Vaccine. **ONLY** use sterile 0.9% Sodium Chloride Injection, USP as the diluent. This diluent is not packaged with the vaccine and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent. Do not add more than 1.8 mL of diluent.


After dilution, 1 vial contains 6 doses of 0.3 mL.

Dilution and Preparation Instructions	
Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – VIAL VERIFICATION	
 <p>✓ Purple plastic cap and purple label border.</p>	Verify that the vial of Pfizer-BioNTech COVID-19 Vaccine has a purple plastic cap. Some vials also may have a purple label border.

Dilution and Preparation Instructions	
Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – THAWING PRIOR TO DILUTION	
 <p>No more than 2 hours at room temperature (up to 25°C/77°F).</p>	<ul style="list-style-type: none"> Thaw vial(s) of Pfizer-BioNTech COVID-19 Vaccine before use either by: <ul style="list-style-type: none"> Allowing vial(s) to thaw in the refrigerator [2°C to 8°C (35°F to 46°F)]. A carton of vials may take up to 3 hours to thaw, and thawed vials can be stored in the refrigerator for up to 1 month. Allowing vial(s) to sit at room temperature [up to 25°C (77°F)] for 30 minutes. Using either thawing method, vials must reach room temperature before dilution and must be diluted within 2 hours.
 <p>Gently × 10</p>	<ul style="list-style-type: none"> Before dilution invert vaccine vial gently 10 times. <u>Do not shake.</u> Inspect the liquid in the vial prior to dilution. The liquid is a white to off-white suspension and may contain white to off-white opaque amorphous particles. Do not use if liquid is discolored or if other particles are observed.


Dilution and Preparation Instructions

Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – DILUTION



Add 1.8 mL of sterile 0.9% sodium chloride injection, USP.

- Obtain sterile 0.9% Sodium Chloride Injection, USP. Use only this as the diluent.
- Using aseptic technique, withdraw 1.8 mL of diluent into a transfer syringe (21-gauge or narrower needle).
- Cleanse the vaccine vial stopper with a single-use antiseptic swab.
- Add 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP into the vaccine vial.



Pull back plunger to 1.8 mL to remove air from vial.

Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.

Dilution and Preparation Instructions




Gently × 10

- Gently invert the vial containing the Pfizer-BioNTech COVID-19 Vaccine 10 times to mix.
- Do not shake.
- Inspect the vaccine in the vial.
- The vaccine will be an off-white suspension. Do not use if vaccine is discolored or contains particulate matter.



**Record the date and time of dilution.
Use within 6 hours after dilution.**

- Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label.
- Store between 2°C to 25°C (35°F to 77°F).
- Discard any unused vaccine 6 hours after dilution.

Dilution and Preparation Instructions	
Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – WITHDRAWAL OF INDIVIDUAL 0.3 mL DOSES	
 <p>Withdraw 0.3 mL dose of vaccine.</p>	<ul style="list-style-type: none"> Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw <u>0.3 mL</u> of the Pfizer-BioNTech COVID-19 Vaccine preferentially using a low dead-volume syringe and/or needle. Each dose must contain 0.3 mL of vaccine. If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume. Administer immediately.

Administration

Visually inspect each dose in the dosing syringe prior to administration. The vaccine will be an off-white suspension. During the visual inspection,

- verify the final dosing volume of 0.3 mL.
- confirm there are no particulates and that no discoloration is observed.
- do not administer if vaccine is discolored or contains particulate matter.

Administer the Pfizer-BioNTech COVID-19 Vaccine intramuscularly.

After dilution, vials of Pfizer-BioNTech COVID-19 Vaccine with purple caps contain 6 doses of 0.3 mL of vaccine. Low dead-volume syringes and/or needles can be used to extract 6 doses from a single vial. If standard syringes and needles are used, there may not be sufficient volume to extract 6 doses from a single vial.

Irrespective of the type of syringe and needle:

- Each dose must contain 0.3 mL of vaccine.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and content.
- Do not pool excess vaccine from multiple vials.

Contraindications

Do not administer Pfizer-BioNTech COVID-19 Vaccine to individuals with known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Pfizer-BioNTech COVID-19 Vaccine (*see Full EUA Prescribing Information*).

Warnings

Management of Acute Allergic Reactions

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Pfizer-BioNTech COVID-19 Vaccine.

Monitor Pfizer-BioNTech COVID-19 Vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention (CDC) guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

Myocarditis and Pericarditis

Postmarketing data demonstrate increased risks of myocarditis and pericarditis, particularly within 7 days following the second dose. The observed risk is higher among males under 40 years of age than among females and older males. The observed risk is highest in males 12 through 17 years of age. Although some cases required intensive care support, available data from short-term follow-up suggest that most individuals have had resolution of symptoms with conservative management. Information is not yet available about potential long-term sequelae. The CDC has published considerations related to myocarditis and pericarditis after vaccination, including for vaccination of individuals with a history of myocarditis or pericarditis (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/myocarditis.html>).

Syncope

Syncope (fainting) may occur in association with administration of injectable vaccines, in particular in adolescents. Procedures should be in place to avoid injury from fainting.

Altered Immunocompetence

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Pfizer-BioNTech COVID-19 Vaccine.

Limitation of Effectiveness

Pfizer-BioNTech COVID-19 Vaccine may not protect all vaccine recipients.

Adverse Reactions

Adverse Reactions in Clinical Trials

Adverse reactions following administration of the Pfizer-BioNTech COVID-19 Vaccine that have been reported in clinical trials include injection site pain, fatigue, headache, muscle pain, chills, joint pain, fever, injection site swelling, injection site redness, nausea, malaise, lymphadenopathy, decreased appetite, rash, and pain in extremity (see *Full EUA Prescribing Information*).

Adverse Reactions in Post Authorization Experience

Severe allergic reactions, including anaphylaxis, and other hypersensitivity reactions (e.g., rash, pruritus, urticaria, angioedema), diarrhea, vomiting, pain in extremity (arm), and syncope have been reported following administration of the Pfizer-BioNTech COVID-19 Vaccine.

Myocarditis and pericarditis have been reported following administration of the Pfizer-BioNTech COVID-19 Vaccine.

Additional adverse reactions, some of which may be serious, may become apparent with more widespread use of the Pfizer-BioNTech COVID-19 Vaccine.

Use with Other Vaccines

There is no information on the co-administration of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.

INFORMATION TO PROVIDE TO VACCINE RECIPIENTS/CAREGIVERS

As the vaccination provider, you must communicate to the recipient or their caregiver, information consistent with the “Vaccine Information Fact Sheet for Recipients and Caregivers” (and provide a copy or direct the individual to the website www.cvdvaccine.com to obtain the Vaccine Information Fact Sheet) prior to the individual receiving each dose of the Pfizer-BioNTech COVID-19 Vaccine, including:

- FDA has authorized the emergency use of the Pfizer-BioNTech COVID-19 Vaccine, which is not an FDA-approved vaccine.
- The recipient or their caregiver has the option to accept or refuse Pfizer-BioNTech COVID-19 Vaccine.
- The significant known and potential risks and benefits of the Pfizer-BioNTech COVID-19 Vaccine, and the extent to which such risks and benefits are unknown.
- Information about available alternative vaccines and the risks and benefits of those alternatives.

For information on clinical trials that are testing the use of the Pfizer-BioNTech COVID-19 Vaccine to prevent COVID-19, please see www.clinicaltrials.gov.

Provide a vaccination card to the recipient or their caregiver with the date when the recipient needs to return for the second dose of Pfizer-BioNTech COVID-19 Vaccine.

Provide the v-safe information sheet to vaccine recipients/caregivers and encourage vaccine recipients to participate in v-safe. V-safe is a new voluntary smartphone-based tool that uses text messaging and web surveys to check in with people who have been vaccinated to identify potential side effects after COVID-19 vaccination. V-safe asks questions that help CDC monitor the safety of COVID-19 vaccines. V-safe also provides second-dose reminders if needed and live telephone follow-up by CDC if participants report a significant health impact following COVID-19 vaccination. For more information, visit: www.cdc.gov/vsafe.

MANDATORY REQUIREMENTS FOR PFIZER-BIONTECH COVID-19 VACCINE ADMINISTRATION UNDER EMERGENCY USE AUTHORIZATION³

In order to mitigate the risks of using this unapproved product under EUA and to optimize the potential benefit of Pfizer-BioNTech COVID-19 Vaccine, the following items are required. Use of unapproved Pfizer-BioNTech COVID-19 Vaccine for active immunization to prevent COVID-19 under this EUA is limited to the following (all requirements **must** be met):

1. Pfizer-BioNTech COVID-19 Vaccine is authorized for use in individuals 5 years of age and older.
2. The vaccination provider must communicate to the individual receiving the Pfizer-BioNTech COVID-19 Vaccine or their caregiver, information consistent with the “Vaccine Information Fact Sheet for Recipients and Caregivers” prior to the individual receiving Pfizer-BioNTech COVID-19 Vaccine.
3. The vaccination provider must include vaccination information in the state/local jurisdiction’s Immunization Information System (IIS) or other designated system.

³ Vaccination providers administering COMIRNATY (COVID-19 Vaccine, mRNA) must adhere to the same reporting requirements.

4. The vaccination provider is responsible for mandatory reporting of the following to the Vaccine Adverse Event Reporting System (VAERS):
 - vaccine administration errors whether or not associated with an adverse event,
 - serious adverse events* (irrespective of attribution to vaccination),
 - cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and
 - cases of COVID-19 that result in hospitalization or death.

Complete and submit reports to VAERS online at <https://vaers.hhs.gov/reportevent.html>. For further assistance with reporting to VAERS call 1-800-822-7967. The reports should include the words “Pfizer-BioNTech COVID-19 Vaccine EUA” in the description section of the report.

5. The vaccination provider is responsible for responding to FDA requests for information about vaccine administration errors, adverse events, cases of MIS in adults and children, and cases of COVID-19 that result in hospitalization or death following administration of Pfizer-BioNTech COVID-19 Vaccine to recipients.

* Serious adverse events are defined as:

- Death;
- A life-threatening adverse event;
- Inpatient hospitalization or prolongation of existing hospitalization;
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions;
- A congenital anomaly/birth defect;
- An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent 1 of the outcomes listed above.

OTHER ADVERSE EVENT REPORTING TO VAERS AND PFIZER INC.

Vaccination providers may report to VAERS other adverse events that are not required to be reported using the contact information above.


To the extent feasible, report adverse events to Pfizer Inc. using the contact information below or by providing a copy of the VAERS form to Pfizer Inc.

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

ADDITIONAL INFORMATION

For general questions, visit the website or call the telephone number provided below.

To access the most recent Pfizer-BioNTech COVID-19 Vaccine Fact Sheets, please scan the QR code provided below.

Global website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

AVAILABLE ALTERNATIVES

COMIRNATY (COVID-19 Vaccine, mRNA) is an FDA-approved vaccine to prevent COVID-19 caused by SARS-CoV-2. There may be clinical trials or availability under EUA of other COVID-19 vaccines.

COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine intended for individuals 12 years of age and older should not be used for individuals 5 through 11 years of age because of the potential for vaccine administration errors, including dosing errors.

FEDERAL COVID-19 VACCINATION PROGRAM

This vaccine is being made available for emergency use exclusively through the CDC COVID-19 Vaccination Program (the Vaccination Program). Healthcare providers must enroll as providers in the Vaccination Program and comply with the provider requirements. Vaccination providers may not charge any fee for the vaccine and may not charge the vaccine recipient any out-of-pocket charge for administration. However, vaccination providers may seek appropriate reimbursement from a program or plan that covers COVID-19 vaccine administration fees for the vaccine recipient (private insurance, Medicare, Medicaid, Health Resources & Services Administration [HRSA] COVID-19 Uninsured Program for non-insured recipients). For information regarding provider requirements and enrollment in the CDC COVID-19 Vaccination Program, see <https://www.cdc.gov/vaccines/covid-19/provider-enrollment.html>.

Individuals becoming aware of any potential violations of the CDC COVID-19 Vaccination Program requirements are encouraged to report them to the Office of the Inspector General, U.S. Department of Health and Human Services, at 1-800-HHS-TIPS or <https://TIPS.HHS.GOV>.

AUTHORITY FOR ISSUANCE OF THE EUA

The Secretary of Health and Human Services (HHS) has declared a public health emergency that justifies the emergency use of drugs and biological products during the COVID-19 pandemic. In response, FDA has issued an EUA for the unapproved product, Pfizer-BioNTech COVID-19 Vaccine, and for certain uses of FDA-approved COMIRNATY for active immunization against COVID-19.

FDA issued this EUA, based on Pfizer-BioNTech's request and submitted data.

For the authorized uses, although limited scientific information is available, based on the totality of the scientific evidence available to date, it is reasonable to believe that the Pfizer-BioNTech COVID-19 Vaccine and COMIRNATY (COVID-19 Vaccine, mRNA) may be effective for the prevention of COVID-19 in individuals as specified in the *Full EUA Prescribing Information*.

This EUA for the Pfizer-BioNTech COVID-19 Vaccine and COMIRNATY (COVID-19 Vaccine, mRNA) will end when the Secretary of HHS determines that the circumstances justifying the EUA no longer exist or when there is a change in the approval status of the product such that an EUA is no longer needed.

For additional information about Emergency Use Authorization visit FDA at: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>.

The Countermeasures Injury Compensation Program

The Countermeasures Injury Compensation Program (CICP) is a federal program that has been created to help pay for related costs of medical care and other specific expenses to compensate people injured after use of certain medical countermeasures. Medical countermeasures are specific vaccines, medications, devices, or other items used to prevent, diagnose, or treat the public during a public health emergency or a security threat. For more information about CICP regarding the Pfizer-BioNTech COVID-19 Vaccine used to prevent COVID-19, visit www.hrsa.gov/cicp, email cicp@hrsa.gov, or call: 1-855-266-2427.



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1450-17.2

Revised: 09 December 2021

END SHORT VERSION FACT SHEET
Long Version (Full EUA Prescribing Information) Begins On Next Page

FULL EMERGENCY USE AUTHORIZATION (EUA) PRESCRIBING INFORMATION

PFIZER-BIONTECH COVID-19 VACCINE

FULL EMERGENCY USE AUTHORIZATION PRESCRIBING INFORMATION: CONTENTS*

- 1 AUTHORIZED USE**
- 2 DOSAGE AND ADMINISTRATION**
 - 2.1 Preparation for Administration
 - 2.2 Administration Information
 - 2.3 Vaccination Schedule
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
 - 5.1 Management of Acute Allergic Reactions
 - 5.2 Myocarditis and Pericarditis
 - 5.3 Syncope
 - 5.4 Altered Immunocompetence
 - 5.5 Limitation of Effectiveness
- 6 OVERALL SAFETY SUMMARY**
 - 6.1 Clinical Trials Experience
 - 6.2 Post Authorization Experience
- 8 REQUIREMENTS AND INSTRUCTIONS FOR REPORTING
ADVERSE EVENTS AND VACCINE ADMINISTRATION
ERRORS**
- 10 DRUG INTERACTIONS**
- 11 USE IN SPECIFIC POPULATIONS**
 - 11.1 Pregnancy
 - 11.2 Lactation
 - 11.3 Pediatric Use
 - 11.4 Geriatric Use
 - 11.5 Use in Immunocompromised
- 13 DESCRIPTION**
- 14 CLINICAL PHARMACOLOGY**
 - 14.1 Mechanism of Action
- 18 CLINICAL TRIAL RESULTS AND SUPPORTING DATA FOR
EUA**
 - 18.1 Efficacy of Primary Series in Participants 16 Years of Age and Older
 - 18.2 Efficacy of Primary Series in Adolescents 12 Through 15 Years of Age
 - 18.3 Immunogenicity of Primary Series in Adolescents 12 Through 15 Years of Age
 - 18.4 Immunogenicity of a Booster Dose Following a Pfizer-BioNTech COVID-19 Vaccine Primary Series in Participants 18 Through 55 Years of Age
 - 18.5 Immunogenicity in Solid Organ Transplant Recipients
 - 18.6 Immunogenicity of a Booster Dose Following Primary Vaccination with Another Authorized COVID-19 Vaccine
- 19 HOW SUPPLIED/STORAGE AND HANDLING**
- 20 PATIENT COUNSELING INFORMATION**
- 21 CONTACT INFORMATION**

* Sections or subsections omitted from the full emergency use authorization prescribing information are not listed.

FULL EMERGENCY USE AUTHORIZATION (EUA) PRESCRIBING INFORMATION

1 AUTHORIZED USE

Pfizer-BioNTech COVID-19 Vaccine is authorized for use under an Emergency Use Authorization (EUA) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 5 years of age and older.

This EUA Prescribing Information pertains only to Pfizer-BioNTech COVID-19 Vaccine supplied in a multiple dose vial with a purple cap, which is authorized for use in individuals 12 years of age and older.

2 DOSAGE AND ADMINISTRATION

For intramuscular injection only.

The storage, preparation, and administration information in this Prescribing Information apply to the Pfizer-BioNTech COVID-19 Vaccine for individuals 12 years of age and older, which is supplied in a multiple dose vial with a purple cap and **MUST BE DILUTED** before use.

Pfizer-BioNTech COVID-19 Vaccine, Multiple Dose Vial with Purple Cap

Age Range	Dilution Information	Doses Per Vial After Dilution	Dose Volume
12 years and older	Dilute with 1.8 mL sterile 0.9% Sodium Chloride Injection, USP prior to use	6	0.3 mL

2.1 Preparation for Administration

Dose Preparation

Each vial **MUST BE DILUTED** before administering the vaccine.

Prior to Dilution

- The Pfizer-BioNTech COVID-19 Vaccine multiple dose vial with a purple cap contains a volume of 0.45 mL and is supplied as a frozen suspension that does not contain preservative.
- Each vial must be thawed before dilution.
- Vials may be thawed in the refrigerator [2°C to 8°C (35°F to 46°F)] or at room temperature [up to 25°C (77°F)] [*see How Supplied/Storage and Handling (19)*].
- Refer to thawing instructions in the panels below.

Dilution

- Dilute the vial contents using 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP (not provided) to form the Pfizer-BioNTech COVID-19 Vaccine. Do not add more than 1.8 mL of diluent.
- ONLY use sterile 0.9% Sodium Chloride Injection, USP as the diluent. This diluent is not packaged with the vaccine and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent.
- After dilution, 1 vial contains 6 doses of 0.3 mL.

Dilution and Preparation Instructions

Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – VIAL VERIFICATION



✓ Purple plastic cap and purple label border.

Verify that the vial of Pfizer-BioNTech COVID-19 Vaccine has a purple plastic cap. Some vials also may have a purple label border on the label.

Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – THAWING PRIOR TO DILUTION



**No more than
2 hours at room
temperature (up
to 25°C/77°F).**

- Thaw vial(s) of Pfizer-BioNTech COVID-19 Vaccine before use either by:
 - Allowing vial(s) to thaw in the refrigerator [2°C to 8°C (35°F to 46°F)]. A carton of vials may take up to 3 hours to thaw, and thawed vials can be stored in the refrigerator for up to 1 month.
 - Allowing vial(s) to sit at room temperature [up to 25°C (77°F)] for 30 minutes.
- Using either thawing method, vials must reach room temperature before dilution and must be diluted within 2 hours.

Dilution and Preparation Instructions




Gently × 10

- Before dilution invert vaccine vial gently 10 times.
- Do not shake.
- Inspect the liquid in the vial prior to dilution. The liquid is a white to off-white suspension and may contain white to off-white opaque amorphous particles.
- Do not use if liquid is discolored or if other particles are observed.


Dilution and Preparation Instructions

Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – DILUTION



Add 1.8 mL of sterile 0.9% sodium chloride injection, USP.


- Obtain sterile 0.9% Sodium Chloride Injection, USP. Use only this as the diluent.
- Using aseptic technique, withdraw 1.8 mL of diluent into a transfer syringe (21-gauge or narrower needle).
- Cleanse the vaccine vial stopper with a single-use antiseptic swab.
- Add 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP into the vaccine vial.



Pull back plunger to 1.8 mL to remove air from vial.

Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.

Dilution and Preparation Instructions



Gently × 10

- Gently invert the vial containing the Pfizer-BioNTech COVID-19 Vaccine 10 times to mix.
- Do not shake.
- Inspect the vaccine in the vial.
- The vaccine will be an off-white suspension. Do not use if vaccine is discolored or contains particulate matter.




**Record the date and time of dilution.
Use within 6 hours after dilution.**

- Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label.
- Store between 2°C to 25°C (35°F to 77°F).
- Discard any unused vaccine 6 hours after dilution.

Dilution and Preparation Instructions

Pfizer-BioNTech COVID-19 Vaccine Vial with Purple Cap – WITHDRAWAL OF INDIVIDUAL 0.3 mL DOSES



**Withdraw 0.3 mL dose of
vaccine.**

- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw 0.3 mL of the Pfizer-BioNTech COVID-19 Vaccine preferentially using a low dead-volume syringe and/or needle.
- Each dose must contain 0.3 mL of vaccine.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume.
- Administer immediately.

2.2 Administration Information

Visually inspect each dose in the dosing syringe prior to administration. The vaccine will be an off-white suspension. During the visual inspection,

- verify the final dosing volume of 0.3 mL.
- confirm there are no particulates and that no discoloration is observed.
- do not administer if vaccine is discolored or contains particulate matter.

Administer the Pfizer-BioNTech COVID-19 Vaccine intramuscularly.

After dilution, vials of Pfizer-BioNTech COVID-19 Vaccine with purple caps contain 6 doses of 0.3 mL of vaccine. Low dead-volume syringes and/or needles can be used to extract 6 doses from a single vial. If standard syringes and needles are used, there may not be sufficient volume to extract 6 doses from a single vial.

Irrespective of the type of syringe and needle:

- Each dose must contain 0.3 mL of vaccine.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume.
- Do not pool excess vaccine from multiple vials.

2.3 Vaccination Schedule

Primary Series

The Pfizer-BioNTech COVID-19 Vaccine is administered intramuscularly as a primary series of 2 doses (0.3 mL each) 3 weeks apart in individuals 12 years of age and older.

A third primary series dose of the Pfizer-BioNTech COVID-19 Vaccine (0.3 mL) at least 28 days following the second dose is authorized for administration to individuals at least 12 years of age who have undergone solid

organ transplantation, or who are diagnosed with conditions that are considered to have an equivalent level of immunocompromise.

Booster Dose

A single Pfizer-BioNTech COVID-19 Vaccine booster dose (0.3 mL) may be administered at least 6 months after completing a primary series of the Pfizer-BioNTech COVID-19 Vaccine or COMIRNATY to individuals 16 years of age and older.

A single booster dose of the Pfizer-BioNTech COVID-19 Vaccine may be administered to individuals 18 years of age and older as a heterologous booster dose following completion of primary vaccination with another authorized COVID-19 vaccine. The dosing interval for the heterologous booster dose is the same as that authorized for a booster dose of the vaccine used for primary vaccination.

The FDA-approved COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine for individuals 12 years of age and older when prepared according to their respective instructions for use, can be used interchangeably.

COMIRNATY (COVID-19 Vaccine, mRNA) and the two EUA-authorized formulations of Pfizer-BioNTech COVID-19 Vaccine intended for individuals 12 years of age and older should not be used for individuals 5 through 11 years of age because of the potential for vaccine administration errors, including dosing errors.

3 DOSAGE FORMS AND STRENGTHS

Pfizer-BioNTech COVID-19 Vaccine is a suspension for injection.

After preparation, each dose of the Pfizer-BioNTech COVID-19 Vaccine supplied in vials with purple caps is 0.3 mL for individuals 12 years of age and older [*see Dosage and Administration (2.1)*].

4 CONTRAINDICATIONS

Do not administer Pfizer-BioNTech COVID-19 Vaccine to individuals with known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Pfizer-BioNTech COVID-19 Vaccine [*see Description (13)*].

5 WARNINGS AND PRECAUTIONS

5.1 Management of Acute Allergic Reactions

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Pfizer-BioNTech COVID-19 Vaccine.

Monitor Pfizer-BioNTech COVID-19 Vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention (CDC) guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

5.2 Myocarditis and Pericarditis

Postmarketing data demonstrate increased risks of myocarditis and pericarditis, particularly within 7 days following the second dose. The observed risk is higher among males under 40 years of age than among females and older males. The observed risk is highest in males 12 through 17 years of age. Although some cases required intensive care support, available data from short-term follow-up suggest that most individuals have had resolution of symptoms with conservative management. Information is not yet available about potential long-term sequelae. The CDC has published considerations related to myocarditis and pericarditis after vaccination, including for vaccination of individuals with a history of myocarditis or pericarditis (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/myocarditis.html>).

5.3 Syncope

Syncope (fainting) may occur in association with administration of injectable vaccines, in particular in adolescents. Procedures should be in place to avoid injury from fainting.

5.4 Altered Immunocompetence

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Pfizer-BioNTech COVID-19 Vaccine.

5.5 Limitation of Effectiveness

The Pfizer-BioNTech COVID-19 Vaccine may not protect all vaccine recipients.

6 OVERALL SAFETY SUMMARY

It is MANDATORY for vaccination providers to report to the Vaccine Adverse Event Reporting System (VAERS) all vaccine administration errors, all serious adverse events, cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and hospitalized or fatal cases of COVID-19 following vaccination with the Pfizer-BioNTech COVID-19 Vaccine.⁴ To the extent feasible, provide a copy of the VAERS form to Pfizer Inc. Please see the REQUIREMENTS AND INSTRUCTIONS FOR REPORTING ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS section for details on reporting to VAERS and Pfizer Inc.

Primary Series

In clinical studies of participants 16 years of age and older who received Pfizer-BioNTech COVID-19 Vaccine containing 30 mcg of a nucleoside-modified messenger RNA encoding the viral spike (S) glycoprotein of SARS-CoV-2 (30 mcg modRNA), adverse reactions following administration of the primary series included pain at the injection site (84.1%), fatigue (62.9%), headache (55.1%), muscle pain (38.3%), chills (31.9%), joint pain (23.6%), fever (14.2%), injection site swelling (10.5%), injection site redness (9.5%), nausea (1.1%), malaise (0.5%), and lymphadenopathy (0.3%).

In a clinical study in adolescents 12 through 15 years of age who received Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA), adverse reactions following administration of the primary series included pain at the injection site (90.5%), fatigue (77.5%), headache (75.5%), chills (49.2%), muscle pain (42.2%), fever (24.3%), joint pain (20.2%), injection site swelling (9.2%), injection site redness (8.6%), lymphadenopathy (0.8%), and nausea (0.4%).

⁴ Vaccination providers administering COMIRNATY (COVID-19 Vaccine, mRNA) must adhere to the same reporting requirements.

Booster Dose

In a clinical study of participants 18 through 55 years of age, adverse reactions following administration of a booster dose were pain at the injection site (83.0%), fatigue (63.7%), headache (48.4%), muscle pain (39.1%), chills (29.1%), joint pain (25.3%), lymphadenopathy (5.2%), nausea (0.7%), decreased appetite (0.3%), rash (0.3%), and pain in extremity (0.3%).

Post Authorization Experience

Severe allergic reactions, including anaphylaxis, have been reported following administration of the Pfizer-BioNTech COVID-19 Vaccine.

Myocarditis and pericarditis have been reported following administration of the Pfizer-BioNTech COVID-19 Vaccine.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Primary Series

The safety of the primary series Pfizer-BioNTech COVID-19 Vaccine was evaluated in participants 12 years of age and older in two clinical studies conducted in the United States, Europe, Turkey, South Africa, and South America.

Study BNT162-01 (Study 1) was a Phase 1/2, 2-part, dose-escalation trial that enrolled 60 participants, 18 through 55 years of age. Study C4591001 (Study 2) is a Phase 1/2/3, multicenter, multinational, randomized, saline placebo-controlled, observer-blind, dose-finding, vaccine candidate-selection (Phase 1) and efficacy (Phase 2/3) study that has enrolled approximately 46,000 participants, 12 years of age or older. Of these, approximately 43,448 participants [21,720 Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA) encoding the viral spike (S) glycoprotein of SARS-CoV-2; 21,728 placebo] in Phase 2/3 are 16 years of age or older (including 138 and 145 adolescents 16 and 17 years of age in the vaccine and placebo groups, respectively) and 2,260 adolescents are 12 through 15 years of age (1,131 and 1,129 in the vaccine and placebo groups, respectively).

In Study 2, all participants 12 through 15 years of age, and 16 years of age and older in the reactogenicity subset, were monitored for solicited local and systemic reactions and use of antipyretic medication after each vaccination in an electronic diary. Participants are being monitored for unsolicited adverse events, including serious adverse events, throughout the study [from Dose 1 through 1 month (all unsolicited adverse events) or 6 months (serious adverse events) after the last vaccination]. Tables 1 through 6 present the frequency and severity of solicited local and systemic reactions, respectively, within 7 days following each dose of Pfizer-BioNTech COVID 19 Vaccine and placebo.

Participants 16 Years of Age and Older

At the time of the analysis of Study 2 for the EUA, 37,586 [18,801 Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA) and 18,785 placebo] participants 16 years of age or older had been followed for a median of 2 months after the second dose.

The safety evaluation in Study 2 is ongoing. The safety population includes participants 16 years of age and older enrolled by October 9, 2020, and includes safety data accrued through November 14, 2020.

Demographic characteristics in Study 2 were generally similar with regard to age, gender, race, and ethnicity among participants who received Pfizer-BioNTech COVID-19 Vaccine and those who received placebo. Overall, among the total participants who received either the Pfizer-BioNTech COVID-19 Vaccine or placebo, 50.6% were male and 49.4% were female, 83.1% were White, 9.1% were Black or African American, 28.0% were Hispanic/Latino, 4.3% were Asian, and 0.5% were American Indian/Alaska Native.

Solicited Local and Systemic Adverse Reactions

Across both age groups, 18 through 55 years of age and 56 years of age and older, the mean duration of pain at the injection site after Dose 2 was 2.5 days (range 1 to 36 days), for redness 2.6 days (range 1 to 34 days), and for swelling 2.3 days (range 1 to 34 days) for participants in the Pfizer-BioNTech COVID-19 Vaccine group.

Solicited reactogenicity data in 16 and 17 year-old participants are limited.

Table 1: Study 2 – Frequency and Percentages of Participants with Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18 Through 55 Years of Age[‡] – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=2291 n^b (%)	Placebo Dose 1 N^a=2298 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=2098 n^b (%)	Placebo Dose 2 N^a=2103 n^b (%)
Redness^c				
Any (>2 cm)	104 (4.5)	26 (1.1)	123 (5.9)	14 (0.7)
Mild	70 (3.1)	16 (0.7)	73 (3.5)	8 (0.4)
Moderate	28 (1.2)	6 (0.3)	40 (1.9)	6 (0.3)
Severe	6 (0.3)	4 (0.2)	10 (0.5)	0 (0.0)
Swelling^c				
Any (>2 cm)	132 (5.8)	11 (0.5)	132 (6.3)	5 (0.2)
Mild	88 (3.8)	3 (0.1)	80 (3.8)	3 (0.1)
Moderate	39 (1.7)	5 (0.2)	45 (2.1)	2 (0.1)
Severe	5 (0.2)	3 (0.1)	7 (0.3)	0 (0.0)
Pain at the injection site^d				
Any	1904 (83.1)	322 (14.0)	1632 (77.8)	245 (11.7)
Mild	1170 (51.1)	308 (13.4)	1039 (49.5)	225 (10.7)
Moderate	710 (31.0)	12 (0.5)	568 (27.1)	20 (1.0)
Severe	24 (1.0)	2 (0.1)	25 (1.2)	0 (0.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after vaccination.

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

[‡] Eight participants were between 16 and 17 years of age.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

[†] Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

Table 2: Study 2 – Frequency and Percentages of Participants with Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18 Through 55 Years of Age[‡] – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=2291 n^b (%)	Placebo Dose 1 N^a=2298 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=2098 n^b (%)	Placebo Dose 2 N^a=2103 n^b (%)
Fever				
≥38.0°C	85 (3.7)	20 (0.9)	331 (15.8)	10 (0.5)
≥38.0°C to 38.4°C	64 (2.8)	10 (0.4)	194 (9.2)	5 (0.2)
>38.4°C to 38.9°C	15 (0.7)	5 (0.2)	110 (5.2)	3 (0.1)
>38.9°C to 40.0°C	6 (0.3)	3 (0.1)	26 (1.2)	2 (0.1)
>40.0°C	0 (0.0)	2 (0.1)	1 (0.0)	0 (0.0)
Fatigue^c				
Any	1085 (47.4)	767 (33.4)	1247 (59.4)	479 (22.8)
Mild	597 (26.1)	467 (20.3)	442 (21.1)	248 (11.8)
Moderate	455 (19.9)	289 (12.6)	708 (33.7)	217 (10.3)
Severe	33 (1.4)	11 (0.5)	97 (4.6)	14 (0.7)
Headache^c				
Any	959 (41.9)	775 (33.7)	1085 (51.7)	506 (24.1)
Mild	628 (27.4)	505 (22.0)	538 (25.6)	321 (15.3)
Moderate	308 (13.4)	251 (10.9)	480 (22.9)	170 (8.1)
Severe	23 (1.0)	19 (0.8)	67 (3.2)	15 (0.7)
Chills^c				
Any	321 (14.0)	146 (6.4)	737 (35.1)	79 (3.8)
Mild	230 (10.0)	111 (4.8)	359 (17.1)	65 (3.1)
Moderate	82 (3.6)	33 (1.4)	333 (15.9)	14 (0.7)
Severe	9 (0.4)	2 (0.1)	45 (2.1)	0 (0.0)
Vomiting^d				
Any	28 (1.2)	28 (1.2)	40 (1.9)	25 (1.2)
Mild	24 (1.0)	22 (1.0)	28 (1.3)	16 (0.8)
Moderate	4 (0.2)	5 (0.2)	8 (0.4)	9 (0.4)
Severe	0 (0.0)	1 (0.0)	4 (0.2)	0 (0.0)
Diarrhea^c				
Any	255 (11.1)	270 (11.7)	219 (10.4)	177 (8.4)
Mild	206 (9.0)	217 (9.4)	179 (8.5)	144 (6.8)
Moderate	46 (2.0)	52 (2.3)	36 (1.7)	32 (1.5)
Severe	3 (0.1)	1 (0.0)	4 (0.2)	1 (0.0)
New or worsened muscle pain^c				
Any	487 (21.3)	249 (10.8)	783 (37.3)	173 (8.2)
Mild	256 (11.2)	175 (7.6)	326 (15.5)	111 (5.3)
Moderate	218 (9.5)	72 (3.1)	410 (19.5)	59 (2.8)
Severe	13 (0.6)	2 (0.1)	47 (2.2)	3 (0.1)

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=2291 n^b (%)	Placebo Dose 1 N^a=2298 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=2098 n^b (%)	Placebo Dose 2 N^a=2103 n^b (%)
New or worsened joint pain^c				
Any	251 (11.0)	138 (6.0)	459 (21.9)	109 (5.2)
Mild	147 (6.4)	95 (4.1)	205 (9.8)	54 (2.6)
Moderate	99 (4.3)	43 (1.9)	234 (11.2)	51 (2.4)
Severe	5 (0.2)	0 (0.0)	20 (1.0)	4 (0.2)
Use of antipyretic or pain medication ^f	638 (27.8)	332 (14.4)	945 (45.0)	266 (12.6)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

f. Severity was not collected for use of antipyretic or pain medication.

‡ Eight participants were between 16 and 17 years of age.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

Table 3: Study 2 – Frequency and Percentages of Participants with Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Redness^c				
Any (>2 cm)	85 (4.7)	19 (1.1)	120 (7.2)	12 (0.7)
Mild	55 (3.1)	12 (0.7)	59 (3.6)	8 (0.5)
Moderate	27 (1.5)	5 (0.3)	53 (3.2)	3 (0.2)
Severe	3 (0.2)	2 (0.1)	8 (0.5)	1 (0.1)
Swelling^c				
Any (>2 cm)	118 (6.5)	21 (1.2)	124 (7.5)	11 (0.7)
Mild	71 (3.9)	10 (0.6)	68 (4.1)	5 (0.3)
Moderate	45 (2.5)	11 (0.6)	53 (3.2)	5 (0.3)
Severe	2 (0.1)	0 (0.0)	3 (0.2)	1 (0.1)

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Pain at the injection site^d				
Any (>2 cm)	1282 (71.1)	166 (9.3)	1098 (66.1)	127 (7.7)
Mild	1008 (55.9)	160 (8.9)	792 (47.7)	125 (7.6)
Moderate	270 (15.0)	6 (0.3)	298 (18.0)	2 (0.1)
Severe	4 (0.2)	0 (0.0)	8 (0.5)	0 (0.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after vaccination.

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

Table 4: Study 2 – Frequency and Percentages of Participants with Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Fever				
≥38.0°C	26 (1.4)	7 (0.4)	181 (10.9)	4 (0.2)
≥38.0°C to 38.4°C	23 (1.3)	2 (0.1)	131 (7.9)	2 (0.1)
>38.4°C to 38.9°C	1 (0.1)	3 (0.2)	45 (2.7)	1 (0.1)
>38.9°C to 40.0°C	1 (0.1)	2 (0.1)	5 (0.3)	1 (0.1)
>40.0°C	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Fatigue^c				
Any	615 (34.1)	405 (22.6)	839 (50.5)	277 (16.8)
Mild	373 (20.7)	252 (14.1)	351 (21.1)	161 (9.8)
Moderate	240 (13.3)	150 (8.4)	442 (26.6)	114 (6.9)
Severe	2 (0.1)	3 (0.2)	46 (2.8)	2 (0.1)
Headache^c				
Any	454 (25.2)	325 (18.1)	647 (39.0)	229 (13.9)
Mild	348 (19.3)	242 (13.5)	422 (25.4)	165 (10.0)
Moderate	104 (5.8)	80 (4.5)	216 (13.0)	60 (3.6)
Severe	2 (0.1)	3 (0.2)	9 (0.5)	4 (0.2)
Chills^c				
Any	113 (6.3)	57 (3.2)	377 (22.7)	46 (2.8)
Mild	87 (4.8)	40 (2.2)	199 (12.0)	35 (2.1)
Moderate	26 (1.4)	16 (0.9)	161 (9.7)	11 (0.7)
Severe	0 (0.0)	1 (0.1)	17 (1.0)	0 (0.0)

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Vomiting^d				
Any	9 (0.5)	9 (0.5)	11 (0.7)	5 (0.3)
Mild	8 (0.4)	9 (0.5)	9 (0.5)	5 (0.3)
Moderate	1 (0.1)	0 (0.0)	1 (0.1)	0 (0.0)
Severe	0 (0.0)	0 (0.0)	1 (0.1)	0 (0.0)
Diarrhea^c				
Any	147 (8.2)	118 (6.6)	137 (8.3)	99 (6.0)
Mild	118 (6.5)	100 (5.6)	114 (6.9)	73 (4.4)
Moderate	26 (1.4)	17 (0.9)	21 (1.3)	22 (1.3)
Severe	3 (0.2)	1 (0.1)	2 (0.1)	4 (0.2)
New or worsened muscle pain^c				
Any	251 (13.9)	149 (8.3)	477 (28.7)	87 (5.3)
Mild	168 (9.3)	100 (5.6)	202 (12.2)	57 (3.5)
Moderate	82 (4.6)	46 (2.6)	259 (15.6)	29 (1.8)
Severe	1 (0.1)	3 (0.2)	16 (1.0)	1 (0.1)
New or worsened joint pain^c				
Any	155 (8.6)	109 (6.1)	313 (18.9)	61 (3.7)
Mild	101 (5.6)	68 (3.8)	161 (9.7)	35 (2.1)
Moderate	52 (2.9)	40 (2.2)	145 (8.7)	25 (1.5)
Severe	2 (0.1)	1 (0.1)	7 (0.4)	1 (0.1)
Use of antipyretic or pain medication	358 (19.9)	213 (11.9)	625 (37.7)	161 (9.8)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

From an independent report (*Kamar N, Abravanel F, Marion O, et al. Three doses of an mRNA Covid-19 vaccine in solid-organ transplant recipients. N Engl J Med*), in 99 individuals who had undergone various solid organ transplant procedures (heart, kidney, liver, lung, pancreas) 97±8 months previously who received a third vaccine dose, the adverse event profile was similar to that after the second dose and no grade 3 or grade 4 events were reported in recipients who were followed for 1 month following post Dose 3.

Unsolicited Adverse Events

Serious Adverse Events

In Study 2, among participants 16 through 55 years of age who had received at least 1 dose of vaccine or placebo (Pfizer-BioNTech COVID-19 Vaccine = 10,841; placebo = 10,851), serious adverse events from Dose 1 through up to 30 days after Dose 2 in ongoing follow-up were reported by 0.4% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 0.3% of placebo recipients. In a similar analysis, in participants 56 years of age and older (Pfizer-BioNTech COVID-19 Vaccine = 7,960, placebo = 7,934), serious adverse events were

reported by 0.8% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 0.6% of placebo recipients who received at least 1 dose of Pfizer-BioNTech COVID-19 Vaccine or placebo, respectively. In these analyses, 91.6% of study participants had at least 30 days of follow-up after Dose 2.

Appendicitis was reported as a serious adverse event for 12 participants, and numerically higher in the vaccine group, 8 vaccine participants and 4 placebo participants. Currently available information is insufficient to determine a causal relationship with the vaccine. There were no other notable patterns or numerical imbalances between treatment groups for specific categories of serious adverse events (including neurologic, neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Non-Serious Adverse Events

In Study 2 in which 10,841 participants 16 through 55 years of age received Pfizer-BioNTech COVID-19 Vaccine and 10,851 participants received placebo, non-serious adverse events from Dose 1 through up to 30 days after Dose 2 in ongoing follow-up were reported in 29.3% of participants who received Pfizer-BioNTech COVID-19 Vaccine and 13.2% of participants in the placebo group, for participants who received at least 1 dose. Overall in a similar analysis in which 7960 participants 56 years of age and older received Pfizer-BioNTech COVID-19 Vaccine, non-serious adverse events within 30 days were reported in 23.8% of participants who received Pfizer-BioNTech COVID-19 Vaccine and 11.7% of participants in the placebo group, for participants who received at least 1 dose. In these analyses, 91.6% of study participants had at least 30 days of follow-up after Dose 2.

The higher frequency of reported unsolicited non-serious adverse events among Pfizer-BioNTech COVID-19 Vaccine recipients compared to placebo recipients was primarily attributed to local and systemic adverse events reported during the first 7 days following vaccination that are consistent with adverse reactions solicited among participants in the reactogenicity subset and presented in Tables 3 and 4. From Dose 1 through 30 days after Dose 2, reports of lymphadenopathy were imbalanced with notably more cases in the Pfizer-BioNTech COVID-19 Vaccine group (64) vs. the placebo group (6), which is plausibly related to vaccination. Throughout the safety follow-up period to date, Bell's palsy (facial paralysis) was reported by 4 participants in the Pfizer-BioNTech COVID-19 Vaccine group. Onset of facial paralysis was Day 37 after Dose 1 (participant did not receive Dose 2) and Days 3, 9, and 48 after Dose 2. No cases of Bell's palsy were reported in the placebo group. Currently available information is insufficient to determine a causal relationship with the vaccine. There were no other notable patterns or numerical imbalances between treatment groups for specific categories of non-serious adverse events (including other neurologic or neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Adolescents 12 Through 15 Years of Age

In an analysis of Study 2, based on data up to the cutoff date of March 13, 2021, 2,260 adolescents (1,131 Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA); 1,129 placebo) were 12 through 15 years of age. Of these, 1,308 (660 Pfizer-BioNTech COVID-19 Vaccine and 648 placebo) adolescents have been followed for at least 2 months after the second dose. The safety evaluation in Study 2 is ongoing.

Demographic characteristics in Study 2 were generally similar with regard to age, gender, race, and ethnicity among adolescents who received Pfizer-BioNTech COVID-19 Vaccine and those who received placebo. Overall, among the adolescents who received the Pfizer-BioNTech COVID-19 Vaccine, 50.1% were male and 49.9% were female, 85.9% were White, 4.6% were Black or African American, 11.7% were Hispanic/Latino, 6.4% were Asian, and 0.4% were American Indian/Alaska Native.

The mean duration of pain at the injection site after Dose 1 was 2.4 days (range 1 to 10 days), for redness 2.4 days (range 1 to 16 days), and for swelling 1.9 days (range 1 to 5 days) for adolescents in the Pfizer-BioNTech COVID-19 Vaccine group.

Table 5: Study 2 – Frequency and Percentages of Adolescents With Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Adolescents 12 Through 15 Years of Age – Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Redness^c				
Any (>2 cm)	65 (5.8)	12 (1.1)	55 (5.0)	10 (0.9)
Mild	44 (3.9)	11 (1.0)	29 (2.6)	8 (0.7)
Moderate	20 (1.8)	1 (0.1)	26 (2.4)	2 (0.2)
Severe	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Swelling^c				
Any (>2 cm)	78 (6.9)	11 (1.0)	54 (4.9)	6 (0.6)
Mild	55 (4.9)	9 (0.8)	36 (3.3)	4 (0.4)
Moderate	23 (2.0)	2 (0.2)	18 (1.6)	2 (0.2)
Severe	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Pain at the injection site^d				
Any	971 (86.2)	263 (23.3)	866 (78.9)	193 (17.9)
Mild	467 (41.4)	227 (20.1)	466 (42.5)	164 (15.2)
Moderate	493 (43.7)	36 (3.2)	393 (35.8)	29 (2.7)
Severe	11 (1.0)	0 (0.0)	7 (0.6)	0 (0.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after vaccination.

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

Table 6: Study 2 – Frequency and Percentages of Adolescents with Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After Each Dose – Adolescents 12 Through 15 Years of Age – Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Fever				
≥38.0°C	114 (10.1)	12 (1.1)	215 (19.6)	7 (0.6)
≥38.0°C to 38.4°C	74 (6.6)	8 (0.7)	107 (9.8)	5 (0.5)
>38.4°C to 38.9°C	29 (2.6)	2 (0.2)	83 (7.6)	1 (0.1)
>38.9°C to 40.0°C	10 (0.9)	2 (0.2)	25 (2.3)	1 (0.1)

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
>40.0°C	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Fatigue^c				
Any	677 (60.1)	457 (40.6)	726 (66.2)	264 (24.5)
Mild	278 (24.7)	250 (22.2)	232 (21.1)	133 (12.3)
Moderate	384 (34.1)	199 (17.7)	468 (42.7)	127 (11.8)
Severe	15 (1.3)	8 (0.7)	26 (2.4)	4 (0.4)
Headache^c				
Any	623 (55.3)	396 (35.1)	708 (64.5)	263 (24.4)
Mild	361 (32.0)	256 (22.7)	302 (27.5)	169 (15.7)
Moderate	251 (22.3)	131 (11.6)	384 (35.0)	93 (8.6)
Severe	11 (1.0)	9 (0.8)	22 (2.0)	1 (0.1)
Chills^c				
Any	311 (27.6)	109 (9.7)	455 (41.5)	73 (6.8)
Mild	195 (17.3)	82 (7.3)	221 (20.1)	52 (4.8)
Moderate	111 (9.8)	25 (2.2)	214 (19.5)	21 (1.9)
Severe	5 (0.4)	2 (0.2)	20 (1.8)	0 (0.0)
Vomiting^d				
Any	31 (2.8)	10 (0.9)	29 (2.6)	12 (1.1)
Mild	30 (2.7)	8 (0.7)	25 (2.3)	11 (1.0)
Moderate	0 (0.0)	2 (0.2)	4 (0.4)	1 (0.1)
Severe	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Diarrhea^c				
Any	90 (8.0)	82 (7.3)	65 (5.9)	43 (4.0)
Mild	77 (6.8)	72 (6.4)	59 (5.4)	38 (3.5)
Moderate	13 (1.2)	10 (0.9)	6 (0.5)	5 (0.5)
Severe	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
New or worsened muscle pain^c				
Any	272 (24.1)	148 (13.1)	355 (32.4)	90 (8.3)
Mild	125 (11.1)	88 (7.8)	152 (13.9)	51 (4.7)
Moderate	145 (12.9)	60 (5.3)	197 (18.0)	37 (3.4)
Severe	2 (0.2)	0 (0.0)	6 (0.5)	2 (0.2)
New or worsened joint pain^c				
Any	109 (9.7)	77 (6.8)	173 (15.8)	51 (4.7)
Mild	66 (5.9)	50 (4.4)	91 (8.3)	30 (2.8)
Moderate	42 (3.7)	27 (2.4)	78 (7.1)	21 (1.9)
Severe	1 (0.1)	0 (0.0)	4 (0.4)	0 (0.0)

	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine[†] Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Use of antipyretic or pain medication ^f	413 (36.6)	111 (9.8)	557 (50.8)	95 (8.8)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

f. Severity was not collected for use of antipyretic or pain medication.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

Unsolicited Adverse Events

In the following analyses of Study 2 in adolescents 12 through 15 years of age (1,131 of whom received Pfizer-BioNTech COVID-19 Vaccine and 1,129 of whom received placebo), 98.3% of study participants had at least 30 days of follow-up after Dose 2.

Serious Adverse Events

Serious adverse events from Dose 1 through up to 30 days after Dose 2 in ongoing follow-up were reported by 0.4% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 0.1% of placebo recipients. There were no notable patterns or numerical imbalances between treatment groups for specific categories of serious adverse events that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Non-Serious Adverse Events

Non-serious adverse events from Dose 1 through up to 30 days after Dose 2 in ongoing follow-up were reported by 5.8% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 5.8% of placebo recipients. From Dose 1 through 30 days after Dose 2, reports of lymphadenopathy plausibly related to the study intervention were imbalanced, with notably more cases in the Pfizer-BioNTech COVID-19 Vaccine group (7) vs. the placebo group (1). There were no other notable patterns or numerical imbalances between treatment groups for specific categories of non-serious adverse events that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Booster Dose Following a Primary Series of Pfizer-BioNTech COVID-19 Vaccine or COMIRNATY

A subset of Study 2 Phase 2/3 participants of 306 adults 18 through 55 years of age received a booster dose of Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA) approximately 6 months (range of 4.8 to 8.0 months) after completing the primary series. Additionally, a total of 23 Study 2 Phase 1 participants (11 participants 18 through 55 years of age and 12 participants 65 through 85 years of age) received a booster dose of Pfizer-BioNTech COVID-19 Vaccine approximately 8 months (range 7.9 to 8.8 months) after completing the primary series. Safety monitoring after the booster dose was the same as that in the reactogenicity subset who received the primary series.

Among the 306 Phase 2/3 participants, the median age was 42 years (range 19 through 55 years of age), 45.8% were male and 54.2% were female, 81.4% were White, 27.8% were Hispanic/Latino, 9.2% were Black or African American, 5.2% were Asian, and 0.7% were American Indian/Alaska Native. Among the 12 Phase 1

participants 65 through 85 years of age, the median age was 69 years (range 65 through 75 years of age), 6 were male and all were White and Not Hispanic/Latino. Following the booster dose, the median follow-up time was 2.6 months (range 2.1 to 2.9 months) for Phase 1 participants and 2.6 months (range 1.1 to 2.8 months) for Phase 2/3 participants.

Solicited Local and Systemic Adverse Reactions

Table 7 and Table 8 present the frequency and severity of reported solicited local and systemic reactions, respectively, within 7 days of a booster dose of Pfizer-BioNTech COVID-19 Vaccine for Phase 2/3 participants 18 through 55 years of age.

In participants who received a booster dose, the mean duration of pain at the injection site after the booster dose was 2.6 days (range 1 to 8 days), for redness 2.2 days (range 1 to 15 days), and for swelling 2.2 days (range 1 to 8 days).

Table 7: Study 2 – Frequency and Percentages of Participants With Solicited Local Reactions, By Maximum Severity, Within 7 Days After the Booster Dose of Pfizer-BioNTech COVID-19 Vaccine – Participants 18 through 55 Years of Age*

Solicited Local Reaction	Pfizer-BioNTech COVID-19 Vaccine† Booster Dose N^a = 289 n^b (%)
Redness^c	
Any (>2 cm)	17 (5.9)
Mild	10 (3.5)
Moderate	7 (2.4)
Severe	0
Swelling^c	
Any (>2 cm)	23 (8.0)
Mild	13 (4.5)
Moderate	9 (3.1)
Severe	1 (0.3)
Pain at the injection site^d	
Any	240 (83.0)
Mild	174 (60.2)
Moderate	65 (22.5)
Severe	1 (0.3)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after the booster dose.

Note: No Grade 4 solicited local reactions were reported.

* A subset of Phase 2/3 participants 18 through 55 years of age who received a booster dose of COMIRNATY (COVID-19 Vaccine, mRNA) approximately 6 months after completing the primary series.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to 5.0 cm; Moderate: >5.0 to 10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

Table 8: Study 2 – Frequency and Percentages of Participants With Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After the Booster Dose of Pfizer-BioNTech COVID-19 Vaccine – Participants 18 through 55 Years of Age*

Solicited Systemic Reaction	Pfizer-BioNTech COVID-19 Vaccine[†] Booster Dose N^a = 289 n^b (%)
Fever	
≥38.0°C	25 (8.7)
≥38.0°C to 38.4°C	12 (4.2)
>38.4°C to 38.9°C	12 (4.2)
>38.9°C to 40.0°C	1 (0.3)
>40.0°C	0
Fatigue^c	
Any	184 (63.7)
Mild	68 (23.5)
Moderate	103 (35.6)
Severe	13 (4.5)
Headache^c	
Any	140 (48.4)
Mild	83 (28.7)
Moderate	54 (18.7)
Severe	3 (1.0)
Chills^c	
Any	84 (29.1)
Mild	37 (12.8)
Moderate	44 (15.2)
Severe	3 (1.0)
Vomiting^d	
Any	5 (1.7)
Mild	5 (1.7)
Moderate	0
Severe	0
Diarrhea^e	
Any	25 (8.7)
Mild	21 (7.3)
Moderate	4 (1.4)
Severe	0

Pfizer-BioNTech COVID-19 Vaccine[†]	
Booster Dose	
N^a = 289	
n^b (%)	
Solicited Systemic Reaction	
New or worsened muscle pain ^c	
Any	113 (39.1)
Mild	52 (18.0)
Moderate	57 (19.7)
Severe	4 (1.4)
New or worsened joint pain ^c	
Any	73 (25.3)
Mild	36 (12.5)
Moderate	36 (12.5)
Severe	1 (0.3)
Use of antipyretic or pain medication ^f	135 (46.7)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after the booster dose.

Note: No Grade 4 solicited systemic reactions were reported.

* A subset of Phase 2/3 participants 18 through 55 years of age who received a booster dose of COMIRNATY (COVID-19 Vaccine, mRNA) approximately 6 months after completing the primary series.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

f. Severity was not collected for use of antipyretic or pain medication.

In Phase 1 participants ≥ 65 years of age (n = 12), local reaction pain at the injection site (n = 8, 66.7%) and systemic reactions fatigue (n = 5, 41.7%), headache (n = 5, 41.7%), chills (n = 2, 16.7%), muscle pain (n = 4, 33.3%), and joint pain (n = 2, 16.7%) were reported after the booster dose. No participant in this age group reported a severe systemic event or fever after the booster dose.

Unsolicited Adverse Events

Overall, the 306 participants who received a booster dose, had a median follow-up time of 2.6 months after the booster dose to the cut-off date (June 17, 2021).

In an analysis of all unsolicited adverse events reported following the booster dose, through 1 month after the booster dose, in participants 18 through 55 years of age (N = 306), those assessed as adverse reactions not already captured by solicited local and systemic reactions were lymphadenopathy (n = 16, 5.2%), nausea (n = 2, 0.7%), decreased appetite (n = 1, 0.3%), rash (n = 1, 0.3%), and pain in extremity (n = 1, 0.3%).

Serious Adverse Events

Of the 306 participants who received a booster dose of Pfizer-BioNTech COVID-19 Vaccine, there were no serious adverse events reported from the booster dose through 30 days after the booster dose. One participant reported a serious adverse event 61 days after the booster dose that was assessed as unrelated to vaccination.

Booster Dose Following Primary Vaccination with Another Authorized COVID-19 Vaccine

The safety of a Pfizer-BioNTech COVID-19 Vaccine booster dose (30 mcg modRNA) in individuals who completed primary vaccination with another authorized COVID-19 Vaccine (heterologous booster dose) is inferred from the safety of a Pfizer-BioNTech COVID-19 Vaccine booster dose administered following completion of Pfizer-BioNTech COVID-19 Vaccine primary series (homologous booster dose) and from data from an independent National Institutes of Health (NIH) study Phase 1/2 open-label clinical trial (NCT04889209) conducted in the United States that evaluated a heterologous booster dose of the Pfizer-BioNTech COVID-19 Vaccine. In this study, adults who had completed primary vaccination with a Moderna COVID-19 Vaccine 2-dose series (N=151), a Janssen COVID-19 Vaccine single dose (N=156), or a Pfizer-BioNTech COVID-19 Vaccine 2-dose series (N=151) at least 12 weeks prior to enrollment and who reported no history of SARS-CoV-2 infection were randomized 1:1:1 to receive a booster dose of one of three vaccines: Moderna COVID-19 Vaccine, Janssen COVID-19 Vaccine, or Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA). Adverse events were assessed through 28 days after the booster dose. An overall review of adverse reactions reported in the study following the Pfizer-BioNTech COVID-19 Vaccine heterologous booster dose did not identify any new safety concerns, as compared with adverse reactions reported following a Pfizer-BioNTech COVID-19 Vaccine primary series doses or homologous booster dose.

6.2 Post Authorization Experience

The following adverse reactions have been identified during post authorization use of Pfizer-BioNTech COVID-19 Vaccine. Because these reactions are reported voluntarily, it is not always possible to reliably estimate their frequency or establish a causal relationship to vaccine exposure.

Cardiac Disorders: myocarditis, pericarditis

Gastrointestinal Disorders: diarrhea, vomiting

Immune System Disorders: severe allergic reactions, including anaphylaxis, and other hypersensitivity reactions (e.g., rash, pruritus, urticaria, angioedema)

Musculoskeletal and Connective Tissue Disorders: pain in extremity (arm)

Nervous System Disorders: syncope

8 REQUIREMENTS AND INSTRUCTIONS FOR REPORTING ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS⁵

See Overall Safety Summary (Section 6) for additional information.

The vaccination provider enrolled in the federal COVID-19 Vaccination Program is responsible for MANDATORY reporting of the listed events following Pfizer-BioNTech COVID-19 Vaccine to the Vaccine Adverse Event Reporting System (VAERS):

- Vaccine administration errors whether or not associated with an adverse event
- Serious adverse events* (irrespective of attribution to vaccination)
- Cases of Multisystem Inflammatory Syndrome (MIS) in children and adults
- Cases of COVID-19 that result in hospitalization or death

⁵ Vaccination providers administering COMIRNATY (COVID-19 Vaccine, mRNA) must adhere to the same reporting requirements.

*Serious adverse events are defined as:

- Death
- A life-threatening adverse event
- Inpatient hospitalization or prolongation of existing hospitalization
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions
- A congenital anomaly/birth defect
- An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent 1 of the outcomes listed above

Instructions for Reporting to VAERS

The vaccination provider enrolled in the federal COVID-19 Vaccination Program should complete and submit a VAERS form to FDA using 1 of the following methods:

- Complete and submit the report online: <https://vaers.hhs.gov/reportevent.html>, or
- If you are unable to submit this form electronically, you may fax it to VAERS at 1-877-721-0366. If you need additional help submitting a report you may call the VAERS toll-free information line at 1-800-822-7967 or send an email to info@vaers.org.

IMPORTANT: When reporting adverse events or vaccine administration errors to VAERS, please complete the entire form with detailed information. It is important that the information reported to FDA be as detailed and complete as possible. Information to include:

- Patient demographics (e.g., patient name, date of birth)
- Pertinent medical history
- Pertinent details regarding admission and course of illness
- Concomitant medications
- Timing of adverse event(s) in relationship to administration of the Pfizer-BioNTech COVID-19 Vaccine
- Pertinent laboratory and virology information
- Outcome of the event and any additional follow-up information if it is available at the time of the VAERS report. Subsequent reporting of follow-up information should be completed if additional details become available.

The following steps are highlighted to provide the necessary information for safety tracking:

1. In Box 17, provide information on Pfizer-BioNTech COVID-19 Vaccine and any other vaccines administered on the same day; and in Box 22, provide information on any other vaccines received within 1 month prior.
2. In Box 18, description of the event:
 - a. Write “Pfizer-BioNTech COVID-19 Vaccine EUA” as the first line.
 - b. Provide a detailed report of vaccine administration error and/or adverse event. It is important to provide detailed information regarding the patient and adverse event/medication error for ongoing safety evaluation of this unapproved vaccine. Please see information to include listed above.
3. Contact information:
 - a. In Box 13, provide the name and contact information of the prescribing healthcare provider or institutional designee who is responsible for the report.
 - b. In Box 14, provide the name and contact information of the best doctor/healthcare professional to contact about the adverse event.

- c. In Box 15, provide the address of the facility where vaccine was given (NOT the healthcare provider's office address).

Other Reporting Instructions

Vaccination providers may report to VAERS other adverse events that are not required to be reported using the contact information above.

To the extent feasible, report adverse events to Pfizer Inc. using the contact information below or by providing a copy of the VAERS form to Pfizer Inc.

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

10 DRUG INTERACTIONS

There are no data to assess the concomitant administration of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.

11 USE IN SPECIFIC POPULATIONS

11.1 Pregnancy

Risk Summary

All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Available data on Pfizer-BioNTech COVID-19 Vaccine administered to pregnant women are insufficient to inform vaccine-associated risks in pregnancy.

In a reproductive and developmental toxicity study, 0.06 mL of a vaccine formulation containing the same quantity of nucleoside-modified messenger ribonucleic acid (modRNA) (30 mcg) and other ingredients included in a single human dose of Pfizer-BioNTech COVID-19 Vaccine was administered to female rats by the intramuscular route on 4 occasions: 21 and 14 days prior to mating, and on gestation days 9 and 20. No vaccine-related adverse effects on female fertility, fetal development, or postnatal development were reported in the study.

11.2 Lactation

Risk Summary

Data are not available to assess the effects of Pfizer-BioNTech COVID-19 Vaccine on the breastfed infant or on milk production/excretion.

11.3 Pediatric Use

Emergency Use Authorization of this formulation of Pfizer-BioNTech COVID-19 Vaccine, supplied in multiple dose vials with purple caps, in adolescents 12 through 17 years of age is based on safety and effectiveness data in this age group and in adults.

For individuals 5 through 11 years of age, a different formulation of the Pfizer-BioNTech COVID-19 Vaccine is authorized.

Emergency Use Authorization of Pfizer-BioNTech COVID-19 Vaccine does not include use in individuals younger than 5 years of age.

11.4 Geriatric Use

Clinical studies of Pfizer-BioNTech COVID-19 Vaccine include participants 65 years of age and older who received the primary series and their data contributes to the overall assessment of safety and efficacy [*see Overall Safety Summary (6.1) and Clinical Trial Results and Supporting Data for EUA (18.1)*]. Of the total number of Pfizer-BioNTech COVID-19 Vaccine recipients in Study 2 (N=20,033), 21.4% (n=4,294) were 65 years of age and older and 4.3% (n=860) were 75 years of age and older.

The safety of a booster dose of Pfizer-BioNTech COVID-19 Vaccine in individuals 65 years of age and older is based on safety data in 12 booster dose recipients 65 through 85 years of age and 306 booster dose recipients 18 through 55 years of age in Study 2. The effectiveness of a booster dose of Pfizer-BioNTech COVID-19 Vaccine in individuals 65 years of age and older is based on effectiveness data in 306 booster dose recipients 18 through 55 years of age in Study 2.

11.5 Use in Immunocompromised

From an independent report (*Kamar N, Abravanel F, Marion O, et al. Three doses of an mRNA Covid-19 vaccine in solid-organ transplant recipients. N Engl J Med*), safety and effectiveness of a third dose of the Pfizer-BioNTech COVID-19 vaccine have been evaluated in persons that received solid organ transplants. The administration of a third dose of vaccine appears to be only moderately effective in increasing potentially protective antibody titers. Patients should still be counselled to maintain physical precautions to help prevent COVID-19. In addition, close contacts of immunocompromised persons should be vaccinated as appropriate for their health status.

13 DESCRIPTION

The Pfizer-BioNTech COVID-19 Vaccine is supplied as a frozen suspension in multiple dose vials with purple caps; each vial must be diluted with 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP prior to use to form the vaccine. Each 0.3 mL dose of the Pfizer-BioNTech COVID-19 Vaccine supplied in multiple dose vials with purple caps contains 30 mcg of a nucleoside-modified messenger RNA (modRNA) encoding the viral spike (S) glycoprotein of SARS-CoV-2.

Each 0.3 mL dose of the Pfizer-BioNTech COVID-19 Vaccine supplied in multiple dose vials with purple caps also includes the following ingredients: lipids (0.43 mg (4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate), 0.05 mg 2[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide, 0.09 mg 1,2-distearoyl-sn-glycero-3-phosphocholine, and 0.2 mg cholesterol), 0.01 mg potassium chloride, 0.01 mg monobasic potassium phosphate, 0.36 mg sodium chloride, 0.07 mg dibasic sodium phosphate dihydrate, and 6 mg sucrose. The

diluent (sterile 0.9% Sodium Chloride Injection, USP) contributes an additional 2.16 mg sodium chloride per dose.

The Pfizer-BioNTech COVID-19 Vaccine does not contain preservative. The vial stoppers are not made with natural rubber latex.

14 CLINICAL PHARMACOLOGY

14.1 Mechanism of Action

The modRNA in the Pfizer-BioNTech COVID-19 Vaccine is formulated in lipid particles, which enable delivery of the RNA into host cells to allow expression of the SARS-CoV-2 S antigen. The vaccine elicits an immune response to the S antigen, which protects against COVID-19.

18 CLINICAL TRIAL RESULTS AND SUPPORTING DATA FOR EUA

18.1 Efficacy of Primary Series in Participants 16 Years of Age and Older

Study 2 is a multicenter, multinational, Phase 1/2/3, randomized, placebo-controlled, observer-blind, dose-finding, vaccine candidate-selection, and efficacy study in participants 12 years of age and older. Randomization was stratified by age: 12 through 15 years of age, 16 through 55 years of age, or 56 years of age and older, with a minimum of 40% of participants in the ≥ 56 -year stratum. The study excluded participants who were immunocompromised and those who had previous clinical or microbiological diagnosis of COVID-19. Participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, were included as were participants with known stable infection with human immunodeficiency virus (HIV), hepatitis C virus (HCV), or hepatitis B virus (HBV).

In the Phase 2/3 portion of Study 2, based on data accrued through November 14, 2020, approximately 44,000 participants 12 years of age and older were randomized equally and received 2 doses of Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA) or placebo separated by 21 days. Participants are planned to be followed for up to 24 months, for assessments of safety and efficacy against COVID-19.

The population for the analysis of the primary efficacy endpoint included, 36,621 participants 12 years of age and older (18,242 in the Pfizer-BioNTech COVID-19 Vaccine group and 18,379 in the placebo group) who did not have evidence of prior infection with SARS-CoV-2 through 7 days after the second dose. Table 9 presents the specific demographic characteristics in the studied population.

Table 9: Demographics (population for the primary efficacy endpoint)^a

	Pfizer-BioNTech COVID-19 Vaccine* (N=18,242) n (%)	Placebo (N=18,379) n (%)
Sex		
Male	9318 (51.1)	9225 (50.2)
Female	8924 (48.9)	9154 (49.8)

	Pfizer-BioNTech COVID-19 Vaccine* (N=18,242) n (%)	Placebo (N=18,379) n (%)
Age (years)		
Mean (SD)	50.6 (15.70)	50.4 (15.81)
Median	52.0	52.0
Min, max	(12, 89)	(12, 91)
Age group		
≥12 through 15 years ^b	46 (0.3)	42 (0.2)
≥16 through 17 years	66 (0.4)	68 (0.4)
≥16 through 64 years	14,216 (77.9)	14,299 (77.8)
≥65 through 74 years	3176 (17.4)	3226 (17.6)
≥75 years	804 (4.4)	812 (4.4)
Race		
White	15,110 (82.8)	15,301 (83.3)
Black or African American	1617 (8.9)	1617 (8.8)
American Indian or Alaska Native	118 (0.6)	106 (0.6)
Asian	815 (4.5)	810 (4.4)
Native Hawaiian or other Pacific Islander	48 (0.3)	29 (0.2)
Other ^c	534 (2.9)	516 (2.8)
Ethnicity		
Hispanic or Latino	4886 (26.8)	4857 (26.4)
Not Hispanic or Latino	13,253 (72.7)	13,412 (73.0)
Not reported	103 (0.6)	110 (0.6)
Comorbidities^d		
Yes	8432 (46.2)	8450 (46.0)
No	9810 (53.8)	9929 (54.0)

* Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

- All eligible randomized participants who receive all vaccination(s) as randomized within the predefined window, have no other important protocol deviations as determined by the clinician, and have no evidence of SARS-CoV-2 infection prior to 7 days after Dose 2.
- 100 participants 12 through 15 years of age with limited follow-up in the randomized population received at least 1 dose (49 in the vaccine group and 51 in the placebo group). Some of these participants were included in the efficacy evaluation depending on the population analyzed. They contributed to exposure information but with no confirmed COVID-19 cases, and did not affect efficacy conclusions.
- Includes multiracial and not reported.
- Number of participants who have 1 or more comorbidities that increase the risk of severe COVID-19 disease
 - Chronic lung disease (e.g., emphysema and chronic bronchitis, idiopathic pulmonary fibrosis, and cystic fibrosis) or moderate to severe asthma
 - Significant cardiac disease (e.g., heart failure, coronary artery disease, congenital heart disease, cardiomyopathies, and pulmonary hypertension)
 - Obesity (body mass index ≥ 30 kg/m²)
 - Diabetes (Type 1, Type 2 or gestational)
 - Liver disease
 - Human Immunodeficiency Virus (HIV) infection (not included in the efficacy evaluation)

The population in the primary efficacy analysis included all participants 12 years of age and older who had been enrolled from July 27, 2020, and followed for the development of COVID-19 through November 14, 2020. Participants 18 through 55 years of age and 56 years of age and older began enrollment from July 27, 2020, 16 through 17 years of age began enrollment from September 16, 2020, and 12 through 15 years of age began enrollment from October 15, 2020.

The vaccine efficacy information is presented in Table 10.

Table 10: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Age Subgroup – Participants Without Evidence of Infection and Participants With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

First COVID-19 occurrence from 7 days after Dose 2 in participants without evidence of prior SARS-CoV-2 infection*			
Subgroup	Pfizer-BioNTech COVID-19 Vaccine[†] N^a=18,198 Cases n1^b Surveillance Time^c (n2^d)	Placebo N^a=18,325 Cases n1^b Surveillance Time^c (n2^d)	Vaccine Efficacy % (95% CI)
All subjects ^e	8 2.214 (17,411)	162 2.222 (17,511)	95.0 (90.3, 97.6) ^f
16 through 64 years	7 1.706 (13,549)	143 1.710 (13,618)	95.1 (89.6, 98.1) ^g
65 years and older	1 0.508 (3848)	19 0.511 (3880)	94.7 (66.7, 99.9) ^g
First COVID-19 occurrence from 7 days after Dose 2 in participants with or without evidence of prior SARS-CoV-2 infection			
Subgroup	Pfizer-BioNTech COVID-19 Vaccine[†] N^a=19,965 Cases n1^b Surveillance Time^c (n2^d)	Placebo N^a=20,172 Cases n1^b Surveillance Time^c (n2^d)	Vaccine Efficacy % (95% CI)
All subjects ^e	9 2.332 (18,559)	169 2.345 (18,708)	94.6 (89.9, 97.3) ^f
16 through 64 years	8 1.802 (14,501)	150 1.814 (14,627)	94.6 (89.1, 97.7) ^g
65 years and older	1 0.530 (4044)	19 0.532 (4067)	94.7 (66.8, 99.9) ^g

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; vomiting).

* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

a. N = Number of participants in the specified group.

b. n1 = Number of participants meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of participants at risk for the endpoint.

e. No confirmed cases were identified in adolescents 12 through 15 years of age.

f. Credible interval for vaccine efficacy (VE) was calculated using a beta-binomial model with a beta (0.700102, 1) prior for $\theta=r(1-VE)/(1+r(1-VE))$, where r is the ratio of surveillance time in the active vaccine group over that in the placebo group.

g. Confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted to the surveillance time.

18.2 Efficacy of Primary Series in Adolescents 12 Through 15 Years of Age

A descriptive efficacy analysis of Study 2 has been performed in approximately 2,200 adolescents 12 through 15 years of age evaluating confirmed COVID-19 cases accrued up to a data cutoff date of March 13, 2021.

The efficacy information in adolescents 12 through 15 years of age is presented in Table 11.

Table 11: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2: Without Evidence of Infection and With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Blinded Placebo-Controlled Follow-up Period, Adolescents 12 Through 15 Years of Age Evaluable Efficacy (7 Days) Population

First COVID-19 occurrence from 7 days after Dose 2 in adolescents 12 through 15 years of age without evidence of prior SARS-CoV-2 infection*			
	Pfizer-BioNTech COVID-19 Vaccine[†] N^a=1005 Cases n1^b Surveillance Time^c (n2^d)	Placebo N^a=978 Cases n1^b Surveillance Time^c (n2^d)	Vaccine Efficacy % (95% CI^e)
Adolescents 12 through 15 years of age	0 0.154 (1001)	16 0.147 (972)	100.0 (75.3, 100.0)
First COVID-19 occurrence from 7 days after Dose 2 in adolescents 12 through 15 years of age with or without evidence of prior SARS-CoV-2 infection			
	Pfizer-BioNTech COVID-19 Vaccine[†] N^a=1119 Cases n1^b Surveillance Time^c (n2^d)	Placebo N^a=1110 Cases n1^b Surveillance Time^c (n2^d)	Vaccine Efficacy % (95% CI^e)
Adolescents 12 through 15 years of age	0 0.170 (1109)	18 0.163 (1094)	100.0 (78.1, 100.0)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; vomiting).

* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

† Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

a. N = Number of participants in the specified group.

b. n1 = Number of participants meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of participants at risk for the endpoint.

e. Confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted for surveillance time.

18.3 Immunogenicity of Primary Series in Adolescents 12 Through 15 Years of Age

In Study 2, an analysis of SARS-CoV-2 50% neutralizing titers (NT50) 1 month after Dose 2 in a randomly selected subset of participants demonstrated non-inferior immune responses (within 1.5-fold) comparing adolescents 12 through 15 years of age to participants 16 through 25 years of age who had no serological or virological evidence of past SARS-CoV-2 infection up to 1 month after Dose 2 (Table 12).

Table 12: Summary of Geometric Mean Ratio for 50% Neutralizing Titer – Comparison of Adolescents 12 Through 15 Years of Age to Participants 16 Through 25 Years of Age (Immunogenicity Subset) –Participants Without Evidence of Infection up to 1 Month After Dose 2 – Dose 2 Evaluable Immunogenicity Population

		Pfizer-BioNTech COVID-19 Vaccine*		12 Through 15 Years/ 16 Through 25 Years	
		12 Through 15 Years n ^a =190	16 Through 25 Years n ^a =170		
Assay	Time Point ^b	GMT ^c (95% CI ^c)	GMT ^c (95% CI ^c)	GMR ^d (95% CI ^d)	Met Noninferiority Objective ^e (Y/N)
SARS-CoV-2 neutralization assay - NT50 (titer) ^f	1 month after Dose 2	1239.5 (1095.5, 1402.5)	705.1 (621.4, 800.2)	1.76 (1.47, 2.10)	Y

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titer; LLOQ = lower limit of quantitation; NAAT = nucleic-acid amplification test; NT50 = 50% neutralizing titer; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Participants who had no serological or virological evidence (up to 1 month after receipt of the last dose) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit up to 1 month after Dose 2 were included in the analysis.

* Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

- n = Number of participants with valid and determinate assay results for the specified assay at the given dose/sampling time point.
- Protocol-specified timing for blood sample collection.
- GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titers and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to $0.5 \times \text{LLOQ}$.
- GMRs and 2-sided 95% CIs were calculated by exponentiating the mean difference of the logarithms of the titers (Group 1 [12 through 15 years of age] – Group 2 [16 through 25 years of age]) and the corresponding CI (based on the Student t distribution).
- Noninferiority is declared if the lower bound of the 2-sided 95% CI for the GMR is greater than 0.67.
- SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralization Assay. The assay uses a fluorescent reporter virus derived from the USA_WA1/2020 strain and virus neutralization is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralized.

18.4 Immunogenicity of a Booster Dose Following a Pfizer-BioNTech COVID-19 Vaccine Primary Series in Participants 18 Through 55 Years of Age

Effectiveness of a booster dose of Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA) was based on an assessment of 50% neutralizing antibody titers (NT50) against SARS-CoV-2 (USA_WA1/2020). In Study 2, analyses of NT50 1 month after the booster dose compared to 1 month after the primary series in individuals 18 through 55 years of age who had no serological or virological evidence of past SARS-CoV-2 infection up to 1 month after the booster vaccination demonstrated noninferiority for both geometric mean ratio (GMR) and difference in seroresponse rates. Seroresponse for a participant was defined as achieving a ≥ 4 -fold rise in NT50 from baseline (before primary series). These analyses are summarized in Table 13 and Table 14.

Table 13: Geometric Mean 50% Neutralizing Titer (SARS-CoV-2 USA_WA1/2020) – Comparison of 1 Month After Booster Dose to 1 Month After Primary Series – Participants 18 Through 55 Years of Age Without Evidence of Infection up to 1 Month After Booster Dose* – Booster Dose Evaluable Immunogenicity Population[±]

Assay	n^a	1 Month After Booster Dose GMT^b (95% CI^b)	1 Month After Primary Series GMT^b (95% CI^b)	1 Month After Booster Dose/ 1 Month After Primary Series GMR^c (97.5% CI^c)	Met Noninferiority Objective^d (Y/N)
SARS-CoV-2 neutralization assay - NT50 (titer) ^e	212	2466.0 (2202.6, 2760.8)	750.6 (656.2, 858.6)	3.29 (2.77, 3.90)	Y

Abbreviations: CI = confidence interval; GMR = geometric mean ratio; GMT = geometric mean titer; LLOQ = lower limit of quantitation; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; NT50 = 50% neutralizing titer; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; Y/N = yes/no.

Note: Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

- * Participants who had no serological or virological evidence (up to 1 month after receipt of a booster dose of Pfizer-BioNTech COVID-19 Vaccine) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative and SARS-CoV-2 not detected by NAAT [nasal swab]) and had a negative NAAT (nasal swab) at any unscheduled visit up to 1 month after the booster dose were included in the analysis.
- ± All eligible participants who had received 2 doses of Pfizer-BioNTech COVID-19 Vaccine as initially randomized, with Dose 2 received within the predefined window (within 19 to 42 days after Dose 1), received a booster dose of Pfizer-BioNTech COVID-19 Vaccine, had at least 1 valid and determinate immunogenicity result after booster dose from a blood collection within an appropriate window (within 28 to 42 days after the booster dose), and had no other important protocol deviations as determined by the clinician.
 - a. n = Number of participants with valid and determinate assay results at both sampling time points within specified window.
 - b. GMTs and 2-sided 95% CIs were calculated by exponentiating the mean logarithm of the titers and the corresponding CIs (based on the Student t distribution). Assay results below the LLOQ were set to $0.5 \times \text{LLOQ}$.
 - c. GMRs and 2-sided 97.5% CIs were calculated by exponentiating the mean differences in the logarithms of the assay and the corresponding CIs (based on the Student t distribution).
 - d. Noninferiority is declared if the lower bound of the 2-sided 97.5% CI for the GMR is >0.67 and the point estimate of the GMR is ≥ 0.80 .
 - e. SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralization Assay. The assay uses a fluorescent reporter virus derived from the USA_WA1/2020 strain and virus neutralization is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralized.

Table 14: Seroreponse Rate for 50% Neutralizing Titer (SARS-CoV-2 USA_WA1/2020) – Comparison of 1 Month After Booster Dose to 1 Month After Primary Series – Participants 18 Through 55 Years of Age Without Evidence of Infection up to 1 Month After Booster Dose* – Booster Dose Evaluable Immunogenicity Population[±]

Assay	N^a	1 Month After Booster Dose n^b % (95% CI^c)	1 Month After Primary Series n^b % (95% CI^c)	Difference (1 Month After Booster Dose - 1 Month After Primary Series) %^d (97.5% CI^e)	Met Noninferiority Objective^f (Y/N)
SARS-CoV-2 neutralization assay - NT50 (titer) ^g	200	199 99.5 (97.2, 100.0)	196 98.0 (95.0, 99.5)	1.5 (-0.7, 3.7)	Y

Abbreviations: CI = confidence interval; LLOQ = lower limit of quantitation; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; NT50 = 50% neutralizing titer; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; Y/N = yes/no.

Note: Seroreponse is defined as achieving a ≥ 4 -fold rise from baseline (before Dose 1). If the baseline measurement is below the LLOQ, a postvaccination assay result $\geq 4 \times$ LLOQ is considered a seroreponse.

Note: Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA).

- * Participants who had no serological or virological evidence (up to 1 month after receipt of booster vaccination) of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative and SARS-CoV-2 not detected by NAAT [nasal swab]) and had a negative NAAT (nasal swab) at any unscheduled visit up to 1 month after booster vaccination were included in the analysis.
- ± All eligible participants who had received 2 doses of Pfizer-BioNTech COVID-19 Vaccine as initially randomized, with Dose 2 received within the predefined window (within 19 to 42 days after Dose 1), received a booster dose of Pfizer-BioNTech COVID-19 Vaccine, had at least 1 valid and determinate immunogenicity result after booster dose from a blood collection within an appropriate window (within 28 to 42 days after the booster dose), and had no other important protocol deviations as determined by the clinician.
- a. N = number of participants with valid and determinate assay results for the specified assay at baseline, 1 month after Dose 2 and 1 month after the booster dose within specified window. These values are the denominators for the percentage calculations.
- b. n = Number of participants with seroreponse for the given assay at the given dose/sampling time point.
- c. Exact 2-sided CI based on the Clopper and Pearson method.
- d. Difference in proportions, expressed as a percentage (1 month after booster dose – 1 month after Dose 2).
- e. Adjusted Wald 2-sided CI for the difference in proportions, expressed as a percentage.
- f. Noninferiority is declared if the lower bound of the 2-sided 97.5% CI for the percentage difference is $> -10\%$.
- g. SARS-CoV-2 NT50 were determined using the SARS-CoV-2 mNeonGreen Virus Microneutralization Assay. The assay uses a fluorescent reporter virus derived from the USA_WA1/2020 strain and virus neutralization is read on Vero cell monolayers. The sample NT50 is defined as the reciprocal serum dilution at which 50% of the virus is neutralized.

18.5 Immunogenicity in Solid Organ Transplant Recipients

From an independent report (*Kamar N, Abravanel F, Marion O, et al. Three doses of an mRNA Covid-19 vaccine in solid-organ transplant recipients. N Engl J Med*), a single arm study has been conducted in 101 individuals who had undergone various solid organ transplant procedures (heart, kidney, liver, lung, pancreas) 97 \pm 8 months previously. A third dose of the Pfizer-BioNTech COVID-19 vaccine was administered to 99 of these individuals approximately 2 months after they had received a second dose. Among the 59 patients who had been seronegative before the third dose, 26 (44%) were seropositive at 4 weeks after the third dose. All 40 patients who had been seropositive before the third dose were still seropositive 4 weeks later. The prevalence of anti-SARS-CoV-2 antibodies was 68% (67 of 99 patients) 4 weeks after the third dose.

18.6 Immunogenicity of a Booster Dose Following Primary Vaccination with Another Authorized COVID-19 Vaccine

Effectiveness of a Pfizer-BioNTech COVID-19 Vaccine booster dose (30 mcg modRNA) in individuals who completed primary vaccination with another authorized COVID-19 Vaccine (heterologous booster dose) is inferred from immunogenicity data supporting effectiveness of a Pfizer-BioNTech COVID-19 Vaccine booster dose administered following completion of Pfizer-BioNTech COVID-19 Vaccine primary series and from immunogenicity data from an independent NIH study Phase 1/2 open-label clinical trial (NCT04889209) conducted in the United States that evaluated a heterologous booster dose of the Pfizer-BioNTech COVID-19 Vaccine. In this study, adults who had completed primary vaccination with a Moderna COVID-19 Vaccine 2-dose series (N=151), a Janssen COVID-19 Vaccine single dose (N=156), or a Pfizer-BioNTech COVID-19 Vaccine 2-dose series (N=151) at least 12 weeks prior to enrollment and who reported no history of SARS-CoV-2 infection were randomized 1:1:1 to receive a booster dose of one of three vaccines: Moderna COVID-19 Vaccine, Janssen COVID-19 Vaccine, or Pfizer-BioNTech COVID-19 Vaccine (30 mcg modRNA). Neutralizing antibody titers, as measured by a pseudovirus neutralization assay using a lentivirus expressing the SARS-CoV-2 Spike protein with D614G mutation, were assessed on Day 1 prior to administration of the booster dose and on Day 15 after the booster dose. A booster response to the Pfizer-BioNTech COVID-19 Vaccine was demonstrated regardless of primary vaccination.

19 HOW SUPPLIED/STORAGE AND HANDLING

The information in this section applies to the Pfizer-BioNTech COVID-19 Vaccine that is supplied in multiple dose vials with a purple cap. These multiple dose vials are supplied in a carton containing 25 multiple dose vials (NDC 59267-1000-3) or 195 multiple dose vials (NDC 59267-1000-2). After dilution, 1 vial contains 6 doses of 0.3 mL. Vial labels and cartons may state that after dilution, a vial contains 5 doses of 0.3 mL. The information in this Full EUA Prescribing Information regarding the number of doses per vial after dilution supersedes the number of doses stated on vial labels and cartons.

During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.

Do not refreeze thawed vials.

Frozen Vials Prior to Use

Cartons of Pfizer-BioNTech COVID-19 Vaccine multiple dose vials with purple caps arrive in thermal containers with dry ice. Once received, remove the vial cartons immediately from the thermal container and preferably store in an ultra-low temperature freezer between -90°C to -60°C (-130°F to -76°F) until the expiry date printed on the label. This information in the package insert supersedes the storage conditions printed on the vial cartons.

Cartons and vials of Pfizer-BioNTech COVID-19 Vaccine supplied in multiple dose vials with purple caps with an expiry date of August 2021 through February 2022 printed on the label may remain in use for 3 months beyond the printed date as long as approved storage conditions between -90°C to -60°C (-130°F to -76°F) have been maintained. Updated expiry dates are shown below.

<u>Printed Expiry Date</u>		<u>Updated Expiry Date</u>
August 2021	→	November 2021
September 2021	→	December 2021
October 2021	→	January 2022
November 2021	→	February 2022
December 2021	→	March 2022
January 2022	→	April 2022
February 2022	→	May 2022

If not stored between -90°C to -60°C (-130°F to -76°F), vials may be stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks. Vials must be kept frozen and protected from light, in the original cartons, until ready to use. Vials stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks may be returned one time to the recommended storage condition of -90°C to -60°C (-130°F to -76°F). Total cumulative time the vials are stored at -25°C to -15°C (-13°F to 5°F) should be tracked and should not exceed 2 weeks.

If an ultra-low temperature freezer is not available, the thermal container in which the Pfizer-BioNTech COVID-19 Vaccine arrives may be used as temporary storage when consistently re-filled to the top of the container with dry ice. Refer to the re-icing guidelines packed in the original thermal container for instructions regarding the use of the thermal container for temporary storage. The thermal container maintains a temperature range of -90°C to -60°C (-130°F to -76°F). Storage of the vials between -96°C to -60°C (-141°F to -76°F) is not considered an excursion from the recommended storage condition.

Transportation of Frozen Vials

If local redistribution is needed and full cartons containing vials cannot be transported at -90°C to -60°C (-130°F to -76°F), vials may be transported at -25°C to -15°C (-13°F to 5°F). Any hours used for transport at -25°C to -15°C (-13°F to 5°F) count against the 2-week limit for storage at -25°C to -15°C (-13°F to 5°F). Frozen vials transported at -25°C to -15°C (-13°F to 5°F) may be returned one time to the recommended storage condition of -90°C to -60°C (-130°F to -76°F).

Thawed Vials Before Dilution

Thawed Under Refrigeration

Thaw and then store undiluted vials in the refrigerator [2°C to 8°C (35°F to 46°F)] for up to 1 month. A carton of 25 vials or 195 vials may take up to 2 or 3 hours, respectively, to thaw in the refrigerator, whereas a fewer number of vials will thaw in less time.

Thawed at Room Temperature

For immediate use, thaw undiluted vials at room temperature [up to 25°C (77°F)] for 30 minutes. Thawed vials can be handled in room light conditions.

Vials must reach room temperature before dilution.

Undiluted vials may be stored at room temperature for no more than 2 hours.

Transportation of Thawed Vials

Available data support transportation of one or more thawed vials at 2°C to 8°C (35°F to 46°F) for up to 12 hours.

Vials After Dilution

After dilution, store vials between 2°C to 25°C (35°F to 77°F) and use within 6 hours from the time of dilution. During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light. Any vaccine remaining in vials must be discarded after 6 hours. Do not refreeze.


20 PATIENT COUNSELING INFORMATION

Advise the recipient or caregiver to read the Vaccine Information Fact Sheet for Recipients and Caregivers.

The vaccination provider must include vaccination information in the state/local jurisdiction's Immunization Information System (IIS) or other designated system. Advise recipient or caregiver that more information about IISs can be found at: <https://www.cdc.gov/vaccines/programs/iis/about.html>.

21 CONTACT INFORMATION

For general questions, visit the website or call the telephone number provided below.

Website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

This Full EUA Prescribing Information may have been updated. For the most recent Full EUA Prescribing Information, please see www.cvdvaccine.com.



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1457-17.2

Revised: 09 December 2021

Revised: 09 December 2021

To:

From: Karen Kingston

AFFIDAVIT OF FINDINGS

Affiant, Karen Kingston (aka Nekedra Karen Kingston), one of the People, Sui Juris, in this Court of Record am presenting the following claims:

Affiant has found the following evidence that the COVID-19 Vaccines are harmful to the American People while engaging in research:

- The COVID-19 vaccines can cause serious adverse events that can result in permanent injuries, autoimmune diseases, chronic diseases, disabilities, and death; per
 - The FDA's October 22, 2020, [COVID-19 Monitoring Vaccine Safety & Efficacy Presentation](#), Slide 17, Exhibit A
 - The December 3, 2021, VAERS database, there have been 946,461 adverse event reports from the COVID-19 vaccines, including almost 20,000 deaths, over 100,000 hospitalizations, and over 32,000 permanent disabilities, Landing Page, Exhibit B
 - According to a [3-year Harvard Study](#), "*...fewer than 1% of vaccine adverse events are reported (in VAERS),*" Page 6, RESULTS, Exhibit C
- Per page 25 of Pfizer's [Biological License Application \(BLA\)](#), approved on August 23, 2021 (per the FDA approval letter), the Pharmacovigilance Plan clearly states that:
 - Heart inflammation (myocarditis and pericarditis) are known risks of the Pfizer COVID-19 vaccine.
 - Vaccine-associated enhanced disease and vaccine enhanced respiratory disease are potential risks of the Pfizer COVID-19 vaccine.
 - Page 25, Exhibit D

Affiant declares that the paramount duty of government is the safety and protection of the People and their property, therefore, Affiant demands immediate investigation into these matters and any potential funds made from any federal program dealing with Covid to be presented to Affiant as it may be a potential conflict of interest.

VERIFICATION

I hereby declare, certify and state, pursuant to the penalties of perjury under the laws of the United States of America, and by the provisions of 28 USC Section 1746, that all of the above and foregoing representations are true and correct to the best of my knowledge, information and belief.

Executed in San Diego, CA on this day DATED the 15th day of December 2021.



Signature of Affiant

CALIFORNIA JURAT WITH AFFIANT STATEMENT

GOVERNMENT CODE § 8202

- ☒ See Attached Document (Notary to cross out lines 1–6 below)
☐ See Statement Below (Lines 1–6 to be completed only by document signer[s], not Notary)

*Signature of Document Signer No. 1*_____
Signature of Document Signer No. 2 (if any)

A notary public or other officer completing this certificate verifies only the identity of the individual who signed the document to which this certificate is attached, and not the truthfulness, accuracy, or validity of that document.

State of California

County of San Diego

Subscribed and sworn to (or affirmed) before me

on this 15 day of December, 2021,
by Date Month Year(1) Nekedra Karen Kingston(and (2) _____),
*Name(s) of Signer(s)*proved to me on the basis of satisfactory evidence to
be the person(s) who appeared before me.*Place Notary Seal and/or Stamp Above*Signature Jennifer Lee Young
*Signature of Notary Public***OPTIONAL**

Completing this information can deter alteration of the document or
fraudulent reattachment of this form to an unintended document.

Description of Attached DocumentTitle or Type of Document: Affidavit of FindingDocument Date: _____ Number of Pages: 1

Signer(s) Other Than Named Above: _____

EXHIBIT A

Go to Slide 17:

https://www.fda.gov/media/143557/download?fbclid=IwAR1SooRjTDuhBPqM4TiD3O7vYgX4eAp3CCqB7SzCk04CMve_OzgtMNPfNkc

FDA Safety Surveillance of COVID-19 Vaccines : DRAFT Working list of possible adverse event outcomes ***Subject to change***

- Guillain-Barré syndrome
- Acute disseminated encephalomyelitis
- Transverse myelitis
- Encephalitis/myelitis/encephalomyelitis/
meningoencephalitis/meningitis/
encepholopathy
- Convulsions/seizures
- Stroke
- Narcolepsy and cataplexy
- Anaphylaxis
- Acute myocardial infarction
- Myocarditis/pericarditis
- Autoimmune disease
- Deaths
- Pregnancy and birth outcomes
- Other acute demyelinating diseases
- Non-anaphylactic allergic reactions
- Thrombocytopenia
- Disseminated intravascular coagulation
- Venous thromboembolism
- Arthritis and arthralgia/joint pain
- Kawasaki disease
- Multisystem Inflammatory Syndrome
in Children
- Vaccine enhanced disease

EXHIBIT B

Go to <https://openvaers.com/covid-data>



Grant Final Report

Grant ID: R18 HS 017045

**Electronic Support for Public Health–Vaccine Adverse
Event Reporting System (ESP:VAERS)**

Inclusive dates: 12/01/07 - 09/30/10

Principal Investigator:

Lazarus, Ross, MBBS, MPH, MMed, GDCompSci

Team members:

Michael Klompas, MD, MPH

Performing Organization:

Harvard Pilgrim Health Care, Inc.

Project Officer:

Steve Bernstein

Submitted to:

The Agency for Healthcare Research and Quality (AHRQ)

U.S. Department of Health and Human Services

540 Gaither Road

Rockville, MD 20850

www.ahrq.gov

EXHIBIT C

Page 6 Results

<https://digital.ahrq.gov/sites/default/files/docs/publication/r18hs017045-lazarus-final-report-2011.pdf>

Results

Preliminary data were collected from June 2006 through October 2009 on 715,000 patients, and 1.4 million doses (of 45 different vaccines) were given to 376,452 individuals. Of these doses, 35,570 possible reactions (2.6 percent of vaccinations) were identified. This is an average of 890 possible events, an average of 1.3 events per clinician, per month. These data were presented at the 2009 AMIA conference.

In addition, ESP:VAERS investigators participated on a panel to explore the perspective of clinicians, electronic health record (EHR) vendors, the pharmaceutical industry, and the FDA towards systems that use proactive, automated adverse event reporting.

Adverse events from drugs and vaccines are common, but underreported. Although 25% of ambulatory patients experience an adverse drug event, less than 0.3% of all adverse drug events and 1-13% of serious events are reported to the Food and Drug Administration (FDA). Likewise, fewer than 1% of vaccine adverse events are reported. Low reporting rates preclude or slow the identification of “problem” drugs and vaccines that endanger public health. New surveillance methods for drug and vaccine adverse effects are needed. Barriers to reporting include a lack of clinician awareness, uncertainty about when and what to report, as well as the burdens of reporting: reporting is not part of clinicians’ usual workflow, takes time, and is duplicative. Proactive, spontaneous, automated adverse event reporting imbedded within EHRs and other information systems has the potential to speed the identification of problems with new drugs and more careful quantification of the risks of older drugs.

Unfortunately, there was never an opportunity to perform system performance assessments because the necessary CDC contacts were no longer available and the CDC consultants responsible for receiving data were no longer responsive to our multiple requests to proceed with testing and evaluation.

EXHIBIT D

Go to page 25 <https://www.fda.gov/media/151733/download>

Pharmacovigilance Plan (PVP)

The Applicant's proposed pharmacovigilance plan (version 1.1) includes the following important risks and missing information:

- Important identified risks: Anaphylaxis; Myocarditis and Pericarditis
- Important potential risk: Vaccine-Associated Enhanced Disease (VAED), including Vaccine-Associated Enhanced Respiratory Disease (VAERD)
- Missing information: Use in pregnancy and lactation; Vaccine effectiveness; Use in pediatric individuals <12 years of age